Opioid prescriber guide

UnitedHealthcare Community Plan of New Mexico

The programs described in this guide were created to help UnitedHealthcare Community Plan members receive the opioid care and treatment they need in safe and effective ways. We've based our measures on Centers for Disease Control and Prevention (CDC) opioid treatment guidelines to help prevent misuse of short-acting and long-acting opioid medications.

Concurrent Drug Utilization Review (cDUR) program

The cDUR program uses the pharmacy claims processing system to screen all prescriptions at the pointof-service and checks for possible inappropriate drug prescribing and utilization, as well as potentially dangerous medical implications or drug interactions. The program includes communication to the dispensing pharmacy at point-of-service through claims edits and messaging. The pharmacist needs to address the clinical situation at the point of sale before entering appropriate National Council for Prescription Drug Programs (NCPDP) codes to receive an approved claim, unless otherwise stated below.

We require prior authorization for concurrent use of opioids and benzodiazepines.
Point-of-sale alert for concurrent use of opioids and MAT drugs.
Point-of-sale alert for concurrent use of opioids and sedative hypnotics.
Point-of-sale alert for concurrent use of opioids and skeletal muscle relaxants.
 Enhanced point-of-sale alert for concurrent use of opioids and prenatal vitamins and concurrent use of opioids and medications used in pregnancy (e.g., doxylamine/pyridoxine) Doesn't require pharmacist to enter appropriate NCPDP codes to receive approved claim
Alerts to concurrent use of multiple SAOs.
Alerts to concurrent use of multiple LAOs.
 Limits combination opioids plus acetaminophen (APAP) Prevents doses of APAP greater than 4 g per day



Concurrent Drug Utilization Review (cDUR) program (cont.)		
High dose opioids - Recommend pharmacist to offer opioid antagonist	 Enhanced point-of-sale alert for opioid doses more than 50 MME that recommends the pharmacist offer an opioid antagonist Doesn't require pharmacist to enter appropriate NCPDP codes to receive approved claim 	
Retrospective Drug Utilization Review (rDUR) programs These programs analyze claims daily and send communications to prescribers.		
Abused medications DUR program	 Identifies members daily who are getting multiple opioid prescriptions from multiple prescribers and/or filling at multiple pharmacies Also identifies members with chronic early refill attempts, overlapping LAOs, overlapping SAOs, high daily doses of opioids, a diagnosis of opioid overdose with an opioid fill, overlapping opioid and MAT medications, overlapping opioid and a benzodiazepine, overlapping opioid and antipsychotic medication, overlapping opioid, muscle relaxant and benzodiazepine, and overlapping opioid and opioid potentiator Sends patient-specific information to all prescribers with medication fill history in last 4 months 	
Pharmacy lock-in program	 Pharmacy lock-in programs vary by state; however, all include filling of opioids by multiple prescribers at multiple pharmacies as an inclusion criterion Requires selected members to use a single pharmacy and/or prescriber(s) for all of their medications 	
Utilization Management (UM) programs These programs help promote appropriate opioid use, reduce costs and improve member health outcomes.		
Cough and cold products containing opioid components	 Limits quantities per fill of 120 mL (units) and a 30-day maximum quantity of 360 mL (units) Requires prior authorization for members under 18 	



Utilization Management (UM) programs (cont.) These programs help promote appropriate opioid use, reduce costs and improve member health outcomes.	
Cumulative 90 milligram morphine equivalent (MME) limit	 Limits dosage at point-of-sale for all opioid products up to 90 MME Prevents the processing of cumulative opioid doses exceeding the limit We require prior authorization for doses above the preset threshold
LAO prior authorization	 Requires prior authorization and the following: Attestation of appropriate use and monitoring Step through SAO (non-cancer pain); step through preferred LAOs If appropriate, step through neuropathic pain alternatives (non-cancer pain)
New-to-therapy SAO edit	 Point-of-sale limit for opioid naïve members (no opioid claims in last 60 days), and includes a maximum 7-day supply and 50 MME or less per day Requires prior authorization to exceed these quantities
Overdose prevention (naloxone)	Prior authorization isn't required for preferred naloxone products (e.g., generic naloxone injection and Narcan® Nasal Spray).
Transmucosal fentanyl product prior authorization	Requires that prior authorization includes documentation of pain due to cancer and patient is already receiving opioids.
Evidence-based prescribing programs These programs focus on outreach to prescribers.	
Fraud/waste/abuse evaluation	 Analyzes retrospective controlled substance claims Identifies outlier opioid prescribers
Miscellaneous	
Miscellaneous – Drug Enforcement Agency (DEA) license edit	Verifies DEA number or license is active and matches scheduled medication in the claim.
Miscellaneous – Refill-too-soon threshold	Increases the refill-too-soon threshold to 90% for opioids and other Schedule CII-V controlled substances.



Pharmacy prior authorization

Information and forms are available on our **Pharmacy Resources and Physician Administered Drugs** page in the Pharmacy Prior Authorization section.

We're here to help

If you have questions, call **888-362-3368.**

Support for your patients:

- 24/7 substance use helpline at **855-780-5955**, available to our members who are your patients, and their caregivers; staffed by licensed behavioral health providers
- liveandworkwell.com

