

UnitedHealthcare® Commercial and Individual Exchange *Medical Policy*

Preventive Care Services

Policy Number: MP.016.53 Effective Date: April 1, 2025

Instructions for Use

Table of Contents	Page
Application	1
Coverage Rationale	
Frequently Asked Questions	
Definitions	
Applicable Codes	5
Benefit Considerations	
Clinical Evidence	
References	
Policy History/Revision Information	
Instructions for Use	

Related Commercial/Individual Exchange Policies

- Breast Imaging for Screening and Diagnosing Cancer
- Cardiovascular Disease Risk Tests
- Colonoscopy Guidelines
- Consultation Services
- Cytological Examination of Breast Fluids for Cancer Screening or Diagnosis
- Genetic Testing for Hereditary Cancer
- Long-Acting Injectable Antiretroviral Agents for HIV (for Commercial Only)
- Long-Acting Injectable Antiretroviral Agents for HIV (for Individual Exchange Only)
- Magnetic Resonance Imaging (MRI) and Computed Tomography (CT) Scan – Site of Service
- Ocular Photoscreening
- Outpatient Surgical Procedures Site of Service
- Preventive Medicine and Screening
- <u>Preventive Vaccines (Immunizations) (for Commercial Only)</u>
- Preventive Vaccines (Immunizations) (for Individual Exchange Only)
- Screening Colonoscopy Procedures Site of Service

Application

UnitedHealthcare Commercial

This Medical Policy applies to UnitedHealthcare Commercial benefit plans.

UnitedHealthcare Individual Exchange

This Medical Policy applies to Individual Exchange benefit plans in all states.

Coverage Rationale

Indications for Coverage Introduction

UnitedHealthcare covers certain medical services under the preventive care services benefit. The federal Patient Protection and Affordable Care Act (PPACA) requires non-grandfathered health plans to cover certain "recommended preventive services" as identified by PPACA under the preventive care services benefit, without cost sharing to members when provided by network providers. This includes:

• Evidence-based items or services that have in effect a rating of "A" or "B" in the current recommendations of the United States Preventive Services Task Force.

Preventive Care Services
Page 1 of 46
UnitedHealthcare Commercial and Individual Exchange Medical Policy
Effective 04/01/2025

- Immunizations for routine use in children, adolescents and adults that have in effect a recommendation from the *Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention*.
- With respect to infants, children and adolescents, evidence-informed preventive care and screenings provided for in the comprehensive guidelines supported by the *Health Resources and Services Administration*.
- With respect to women, such additional preventive care and screenings as provided for in comprehensive guidelines supported by the *Health Resources and Services Administration*.

Member Cost-Sharing Non-Grandfathered Plans

- Non-grandfathered plans provide coverage for preventive care services with no member cost sharing (i.e., covered at 100% of Allowed Amounts without deductible, coinsurance, or copayment) when services are obtained from a Network provider.
- Under PPACA, services obtained from an out-of-network provider are not required to be covered under a plan's preventive benefit and may be subject to member cost sharing. Refer to the member specific benefit plan document for out-of-network benefit information, if any.

Grandfathered Plans

- Plans that maintain grandfathered status under PPACA are not required by law to provide coverage for these preventive services without member cost sharing; although a grandfathered plan may choose to voluntarily amend its plan document to include these preventive benefits.
- Except where there are state mandates, a grandfathered plan might include member cost sharing, or exclude some of the preventive care services identified under PPACA.
- Refer to the member specific benefit plan document for details on how benefits are covered under a grandfathered plan.

Preventive vs. Diagnostic Services

Certain services can be done for preventive or diagnostic reasons. When a service is performed for the purpose of preventive screening and is appropriately reported, it will be considered under the preventive care services benefit. This includes services directly related to the performance of a covered preventive care service (refer to the <u>Frequently Asked Questions</u> section for additional information.)

Preventive services are those performed on a person who:

- has not had the preventive screening done before and does not have symptoms or other abnormal studies suggesting abnormalities*; or
- has had screening done within the recommended interval with the findings considered normal; or
- has had diagnostic services results that were normal after which the physician recommendation would be for future preventive screening studies using the preventive services intervals.

*In the case of a colonoscopy done as a follow-up to a positive stool-based screening (e.g., FIT, FOBT, and fecal DNA) or direct visualization screening (e.g., sigmoidoscopy or CT colonography), refer to FAQ#4 below.

When a service is done for diagnostic purposes, it will be considered under the applicable non-preventive medical benefit. Diagnostic services are done on a person who:

- had abnormalities found on previous preventive or diagnostic studies that require further diagnostic studies; or
- had abnormalities found on previous preventive or diagnostic studies that would recommend a repeat of the same studies within shortened time intervals from the recommended preventive screening time intervals; or
- had a symptom(s) that required further diagnosis; or
- does not fall within the applicable population for a recommendation or guideline.

Covered Breastfeeding Equipment

Personal-use electric breast pump:

- The purchase of a personal-use electric breast pump (HCPCS code E0603).
 - This benefit is limited to one pump per birth. In the case of a birth resulting in multiple infants, only one breast pump is covered.
 - A breast pump purchase includes the necessary supplies for the pump to operate.
- Replacement breast pump supplies necessary for the personal-use electric breast pump to operate. This includes: standard power adaptor, tubing adaptors, tubing, locking rings, bottles specific to breast pump operation, caps for bottles that are specific to the breast pump, valves, filters, and breast shield and/or splash protector for use with the breast pump.

Breastmilk storage bags (HCPCS code A4287).

Colonoscopies

Colonoscopy – Preventive Care Services Benefit (Without Member Cost-Sharing)

Member cost-sharing for colonoscopy is waived when all of the following apply:

- The patient's age is 45-75 years (ends on 76th birthday) as recommended by the USPSTF; and
- The provider is participating in the network; and
- When billed in accordance with the coding in the Colorectal Cancer Screening row listed in this policy.

Colonoscopy may require a site of service review. Refer to the Medical Policies titled <u>Screening Colonoscopy Procedures</u> <u>— Site of Service</u> and <u>Outpatient Surgical Procedures — Site of Service</u>.

Colonoscopy – Medical Benefit (With Member Cost-Sharing)

Member cost-sharing may apply when a colonoscopy is done in any one of the following scenarios:

- The patient's age is outside of the age recommendation of the USPSTF (age 45-75 years); or
- The provider is non-network; or
- Colonoscopy performed with a shortened time interval outside of the USPSTF recommendations; or
- Colonoscopy performed for diagnostic purposes; or
- Colonoscopy performed for surveillance purposes (e.g., a follow-up colonoscopy performed after identification or removal of a polyp or cancer on a previous colonoscopy); or
- Colonoscopy performed for therapeutic/treatment purposes.

The above colonoscopies may require advanced notification and/or site of service review. Refer to the <u>Colonoscopy</u> <u>Guidelines</u> and the Medical Policy titled <u>Outpatient Surgical Procedures – Site of Service</u>.

Coverage Limitations and Exclusions

- Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- The coverage outlined in this policy does not address certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA. These preventive benefits are administered by the member's pharmacy plan administrator. For details on coverage, refer to the member-specific pharmacy plan administrator.
- A vaccine (immunization) is not covered if it does not meet company vaccine policy requirements for FDA labeling and
 if it does not have explicit ACIP recommendations for routine use published in the Morbidity and Mortality Weekly
 Report (MMWR) of the Centers for Disease Control and Prevention (CDC) and is not listed on the applicable
 immunization schedule of ACIP. (Refer to the Preventive Care Services: Vaccine Codes.)
- Examinations, screenings, testing, or vaccines (immunizations) are not covered when:
 - o required solely for the purposes of career or employment, school or education, sports or camp, travel [including travel vaccines (immunizations)], insurance, marriage, or adoption; or
 - o related to judicial or administrative proceedings or orders; or
 - o conducted for purposes of medical research; or
 - o required to obtain or maintain a license of any type.
- Services that are investigational, experimental, unproven, or not medically necessary are not covered.
- Breastfeeding equipment and supplies not listed above. This includes, but is not limited to:
 - Manual breast pumps and all related equipment and supplies.
 - Hospital-grade breast pumps and all related equipment and supplies.
 - Equipment and supplies not listed in the <u>Covered Breastfeeding Equipment</u> section above, including but not limited to:
 - Batteries, battery-powered adaptors, and battery packs.
 - Electrical power adapters for travel.
 - Bottles which are not specific to breast pump operation. This includes the associated bottle nipples, caps, and lids.
 - Travel bags, and other similar travel or carrying accessories.
 - Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
 - Baby weight scales.
 - Garments or other products that allow hands-free pump operation.

- Breast milk storage accessories such as ice-packs, labels, labeling lids, and other similar products. The breastmilk storage accessories exclusion does not apply to breastmilk storage bags (HCPCS code A4287).
- Nursing bras, bra pads, breast shells, nipple shields, and other similar products.
- Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

Note: Refer to the <u>Indications for Coverage</u> section above for covered breastfeeding equipment.

Frequently Asked Questions (FAQ)

1	Q:	If a polyp is encountered during a preventive screening colonoscopy, are future colonoscopies considered under the preventive care services benefit?				
	A:	No. If a polyp is removed during a preventive screening colonoscopy, future colonoscopies would normally be considered to be diagnostic because the time intervals between future colonoscopies would be shortened.				
2	Q:	Are the related therapeutic services for a preventive colonoscopy covered under the preventive care benefit?				
	A: Yes, related services integral to a colonoscopy are covered under the preventive care services benefit including: pre-operative examination, the associated facility, anesthesia, polyp remonecessary), pathologist and physician fees. However, the preventive benefit does not include operative examination.					
3	Q:	Do any preventive care services require prior-authorization?				
	A:	Certain services require prior-authorization on most benefit plans. This includes, but may not be limited to: BRCA lab screening, computed tomographic colonography (virtual colonoscopy), and screening for lung cancer with low-dose computed tomography.				
4						
	A:	Yes, in this situation, the colonoscopy would be considered under the preventive care services benefit when billed in accordance with the coding in the Colorectal Cancer Screening row listed in this policy.				
5	Q:	For preventive services that have a diagnosis code requirement, does the listed diagnosis code need to be the primary diagnosis on the claim?				
	A:	In general, most preventive services do not require the preventive diagnosis code to be in the primary position. However, certain preventive services do require the diagnosis code to be in the primary position, which include: (1) Chemoprevention of Breast Cancer (Counseling), (2) Genetic Counseling and Evaluation for BRCA Testing, and (3) Prevention of Human Immunodeficiency Virus (HIV) Infection.				
If a woman has an abnormal finding on a preventive screening mammog mammogram was found to be normal, will UnitedHealthcare cover her fu		If a woman has an abnormal finding on a preventive screening mammography and the follow up mammogram was found to be normal, will UnitedHealthcare cover her future mammograms under the preventive care services benefit?				
	A:	Yes, if the member was returned to normal mammography screening protocol, her future mammography screenings would be considered under the preventive care services benefit.				
7	Q:	If a member had elevated cholesterol on a prior preventive screening, are future cholesterol tests considered under the preventive care services benefit?				
	A:	Once the diagnosis has been made, further testing is considered diagnostic rather than preventive. This is true whether or not the member is receiving pharmacotherapy.				
8	Q:	Are the related services for a woman's outpatient sterilization or other contraceptive procedure covered under the preventive care benefit?				
	A:	Related services for a woman's outpatient sterilization or other contraceptive procedure are covered under the preventive care services benefit. This includes associated implantable devices, facility fee, anesthesia, and surgeon/physician fees. If a woman is admitted to an inpatient facility for another reason (e.g., maternity/delivery), and has a sterilization or other contraceptive procedure performed during that admission, the sterilization or other contraceptive procedure fees mare covered under				

		the preventive care services benefit. This includes associated sterilization/ contraception surgical fees, device fees, anesthesia, pathology, and physician fees. However, the facility fees are not covered under the preventive care benefit since the sterilization or other contraceptive procedure is incidental to, and is not the primary reason, for the inpatient admission.
9	Q:	Are blood draws/venipunctures included in the preventive care benefit?
	A:	Yes, blood draws/venipunctures are considered under the preventive benefit if billed for a covered preventive lab service that requires a blood draw.
10	Q:	Is a newly-combined vaccine (a vaccine with several individual vaccines combined into one) covered under preventive care benefits?
	A:	A new vaccine that is pending ACIP recommendations, but is a combination of previously approved individual components, may be eligible under the preventive care benefit.
11	Q:	Are preventive care services affected by other policies?
	A:	Yes, including for example, the Reimbursement Policy titled <u>Preventive Medicine and Screening</u> <u>Policy</u> describes situations which may affect reimbursement of preventive care services.
12	Q:	Are travel vaccines covered under preventive care benefits?
	A:	Benefits for preventive care services include vaccines for routine use in children, adolescents and adults that have in effect a recommendation from ACIP with respect to the individual involved. Vaccines that are specific to travel (e.g., typhoid, yellow fever, cholera, plague, and Japanese encephalitis virus) are excluded from the preventive care services benefit.
13	Q:	Does the preventive care services benefit include prescription or over the counter (OTC) items?
	A:	Refer to the plan's pharmacy benefit plan administrator for details on prescription medications and OTCs available under the plan's preventive benefit.

Definitions

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

Modifier 33: Preventive service; when the primary purpose of the service is the delivery of an evidence based service in accordance with a US Preventive Services Task Force A or B rating in effect and other preventive services identified in preventive services mandates (legislative or regulatory), the service may be identified by adding 33 to the procedure. For separately reported services specifically identified as preventive, the modifier should not be used.

Note: UnitedHealthcare considers the procedures and diagnostic codes and Preventive Benefit Instructions listed in the table below in determining whether preventive care benefits apply. While Modifier 33 may be reported, it is not used in making preventive care benefit determinations.

Acronyms

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HRSA: Health Resources and Services Administration

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply. CPT® is a registered trademark of the American Medical Association.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

A date in this column is when the listed rating was released, not when the benefit is effective.

Abdominal Aortic Aneurysm Screening

Service

USPSTF Rating (Dec. 2019): B

The USPSTF recommends 1-time screening for abdominal aortic aneurysm (AAA) with ultrasonography in men aged 65-75 years who have ever smoked.

Code(s)

Procedure Code(s):

Ultrasound Screening Study for Abdominal Aortic Aneurysm: 76706

Diagnosis Code(s):

F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Age 65 through 75 (ends on 76th birthday).

Requires at least one of the diagnosis codes listed in this row.

Bacteriuria Screening

USPSTF Rating (Sept. 2019): A

The USPSTF recommends screening for asymptomatic bacteriuria using urine culture in pregnant persons.

Procedure Code(s):

81007, 87086, 87088

Diagnosis Code(s):

Pregnancy Diagnosis Codes

Requires a Pregnancy Diagnosis Code.

Chlamydia Infection Screening

USPSTF Rating (Sept. 2021): B

The USPSTF recommends screening for chlamydia in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection.

Notes:

- This recommendation applies to asymptomatic, sexually active adolescents and adults, including pregnant persons.
- Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.

Procedure Code(s):

Chlamydia Infection Screening: 86631, 86632, 87110, 87270, 87320, 87490, 87491, 87492, 87801, 87810

Blood Draw:

36415, 36416

Blood draw codes only apply to lab codes 86631 or 86632

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

Other: Z11.3, Z11.4, Z11.8, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51,

Z72.52, Z72.53

Chlamydia Infection Screening:

Requires a <u>Pregnancy Diagnosis Code</u> **or** one of the Screening diagnosis codes listed in this row.

Blood Draw:

Required to be billed with 86631 or 86632 **and**

- One of the Screening diagnosis codes listed in this row **or**
- With a <u>Pregnancy Diagnosis Code</u>.

Gonorrhea Screening

USPSTF Rating (Sept. 2021): B

The USPSTF recommends screening for gonorrhea in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection.

Notes:

 This recommendation applies to asymptomatic, sexually active

Procedure Code(s):

87590, 87591, 87592, 87801, 87850

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Requires either a <u>Pregnancy Diagnosis</u>
<u>Code</u> or one of the Screening diagnosis
codes listed in this row.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

0	ervic	се					
4	date	in	this	column	is	when	th

ne listed rating was released, not when the benefit is effective.

- adolescents and adults including pregnant persons.
- Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.

Code(s)

Other: Z11.3, Z11.4, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Hepatitis B Virus Infection Screening

Pregnant Women:

USPSTF Rating (July 2019): A

The USPSTF recommends screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit.

Adolescents and Adults at *Increased Risk for Infection:*

USPSTF Rating (Dec. 2020): B

The USPSTF recommends screening for hepatitis B virus (HBV) infection in persons at high risk for infection.

Bright Futures (July 2022):

Bright Futures recommends screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).

Also refer to the Medical Policy titled Hepatitis Screening.

Procedure Code(s):

Hepatitis B Virus Infection Screening: 86704, 86706, 87340, 87341, 87467, G0499

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

Other: Z11.3, Z11.4, Z20.2, Z20.6, Z11.59, Z29.81, Z57.8, Z72.51,

Z72.52, Z72.53

Hepatitis B Virus Infection Screening: Requires a Pregnancy Diagnosis Code

or one of the Screening diagnosis codes listed in this row.

Blood Draw:

Requires one of the listed Hepatitis B Virus Infection Screening procedure codes listed in this row and

- A Pregnancy Diagnosis Code or
- One of the Screening diagnosis codes listed in this row.

Hepatitis C Virus Infection Screening

USPSTF Rating (March 2020): B

The USPSTF recommends screening for hepatitis C virus infection in adults aged 18-79 years.

Bright Futures (March 2021)

Bright Futures recommends screening all individuals ages 18 to 79 years at least once for hepatitis C virus infection (HCV).

Also refer to the Medical Policy titled Hepatitis Screening.

Procedure Code(s):

Hepatitis C Virus Infection Screening: 86803, 86804, G0472

Blood Draw: 36415, 36416

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Hepatitis C Virus Infection Screening: Does not have diagnosis code requirements for the preventive benefit to apply.

Blood Draw:

Requires one of the Hepatitis C Virus Infection Screening procedure codes listed in this row

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults

USPSTF Rating (June 2019): A

The USPSTF recommends that clinicians screen for HIV infection in:

- Adolescents and adults aged 15-65 years. Younger adolescents and older adults who are at increased risk of infection should also be screened.
- All pregnant persons, including those who present in labor or at delivery whose HIV status is unknown.

Note: Bright Futures recommends HIV screening lab work be conducted at least once between ages 15-21 years. Also recommended anytime between ages 11-14 years, when a risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.

Code(s)

Procedure Code(s):

HIV (Human Immunodeficiency Virus) Screening:

86689, 86701, 86702, 86703, 87389, 87390, 87391, 87534, 87535, 87536, 87537, 87538, 87539, 87806, G0432, G0433, G0435, G0475, S3645

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129,

Other: Z11.3, Z11.4, Z11.59, Z11.9, Z20.2, Z20.6, Z22.6, Z22.8, Z22.9, Z29.81, Z72.51, Z72.52, Z72.53
Also see Expanded Women's Preventive Health section.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

No age limits.

HIV – Human Immunodeficiency Virus – Screening:

Requires a <u>Pregnancy Diagnosis Code</u> **or** one of the Screening diagnosis codes listed in this row.

Blood Draw:

Requires **both** of the following:

- One of the listed HIV Screening procedure codes listed in this row and
- One of the Screening diagnosis codes listed in this row or a <u>Pregnancy Diagnosis Code</u>.

RH Incompatibility Screening

USPSTF Rating (Feb. 2004): A

Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.

USPSTF Rating (Feb. 2004): B

Repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.

Procedure Code(s):

RH Incompatibility Screening: 86850, 86901

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy Diagnosis Codes

RH Incompatibility Screening:

Requires a Pregnancy Diagnosis Code.

Blood Draw:

Required to be billed with 86850 or 86901 **and** with a <u>Pregnancy Diagnosis</u> Code.

Preventive Care Services

Page 8 of 46
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Page 8 of 46
Effective 04/01/2025

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Syphilis Screening

Non-Pregnant Adolescents and Adults at Increased Risk:

USPSTF Rating (Sept. 2022): A

The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection (asymptomatic, nonpregnant adolescents and adults who are at increased risk for syphilis infection).

Pregnant Women:

USPSTF Rating (Sept. 2018): A

The USPSTF recommends early screening for syphilis infection in all pregnant women.

Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.

Code(s)

Procedure Code(s):

Syphilis Screening: 0064U, 0065U, 0210U, 86592, 86593, 86780

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

Other: Z11.2, Z11.3, Z11.4, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Syphilis Screening:

Requires a <u>Pregnancy Diagnosis Code</u> **or** one of the Screening diagnosis code listed in this row.

Blood Draw:

Requires **both** of the following:

- One of the listed Syphilis Screening procedure codes listed in this row and
- One of the Screening diagnosis codes listed in this row or a <u>Pregnancy Diagnosis Code</u>.

Genetic Counseling and Evaluation for BRCA Testing; and BRCA Lab Screening

USPSTF Rating (Aug. 2019): B

The USPSTF recommends that primary care clinicians assess women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or who have an ancestry associated with breast cancer susceptibility 1 and 2 (BRCA1/2) gene mutations with an appropriate brief familial risk assessment tool. Women with a positive result on the risk assessment tool should receive genetic counseling and, if indicated after counseling, genetic testing.

Refer to the Medical Policy titled Genetic Testing for Hereditary Cancer.

Genetic Counseling and Evaluation

Procedure Code(s):

Medical Genetics and Genetic Counseling Services:

96041, S0265

Evaluation and Management (Office Visits):

99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, 99385, 99386, 99387, 99395, 99396, 99397, G0463

Diagnosis Code(s):

Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43

BRCA Lab Screening Procedure Code(s):

81162, 81163, 81164, 81165, 81166, 81167, 81212, 81215, 81216, 81217

Blood Draw: 36415, 36416

Genetic Counseling and Evaluation

*Medical Necessity plans require genetic counseling before BRCA Lab Screening.

Requires one of the Genetic Counseling and Evaluation diagnosis codes listed in this row in the primary position.

BRCA Lab Screening

*Prior authorization requirements apply to BRCA lab screening.

Applies to **age 18+** when billed with one of the BRCA Lab Screening diagnosis codes listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	m		

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Diagnosis Code(s):

Family History or Personal History of breast cancer and/or ovarian cancer: Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Blood Draw:

Requires one of the BRCA Lab Screening procedure codes listed in this row **and** one of the BRCA Lab Screening diagnosis codes listed in this row.

Screening for Pre-Diabetes and Type 2 Diabetes

USPSTF Rating (Aug. 2021): B

The USPSTF recommends screening for prediabetes and type 2 diabetes in adults aged 35 to 70 years who have overweight or obesity. Clinicians should offer or refer patients with prediabetes to effective preventive interventions.

Refer to Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors:
Behavioral Counseling Interventions for intensive behavioral counseling interventions.

For additional diabetes screening benefits, also see the *Expanded Women's Preventive Health* section for <u>Screening for Diabetes in Pregnancy</u> and <u>Screening for Diabetes After Pregnancy</u>.

Pre-Diabetes Preventive Interventions

Procedure Code(s):

Medical Nutrition Therapy or Counseling: 97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling: 99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0447, G0473, G9886

Diagnosis Code(s):

R73.03 (prediabetes)

Diabetes Screening Procedure Code(s):

Diabetes Screening:

82947, 82948, 82950, 82951, 82952, 83036

Blood Draw: 36415, 36416

Diagnosis Code(s):

Required Diagnosis Codes (requires at least one):

Z00.00, Z00.01, Z13.1

And one of the following additional diagnosis codes as follows:

Additional Diagnosis Codes (requires at least one):

Overweight:

E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29

Obesity:

E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Pre-Diabetes Preventive Interventions

Limited to age 35-70 years (ends on 71st birthday).

Requires diagnosis code R73.03.

Diabetes Screening

Limited to age 35-70 years (ends on 71st birthday).

Diabetes Screening:

Requires one of the Required Diagnosis Codes listed in this row **and** one of the listed Additional Diagnosis Codes in this row

Blood Draw:

Requires **all** of the following:

- One of the listed Diabetes
 Screening procedure codes listed in this row and
- One of the listed Required Diagnosis Codes and
- One of the listed Additional Diagnosis Codes.

Preventive Benefit Does Not Apply:

If a Diabetes Diagnosis Code is present in any position, the preventive benefit does **not** apply; see the <u>Diabetes</u> <u>Diagnosis Code List</u>.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator

For preventive care medications, refe	r to the pharmacy plan administrator.	
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
Gestational Diabetes	Body Mass Index 30.0 – 39.9: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39 Body Mass Index 40.0 and Over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 See the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy. See the Expanded Women's	See the Expanded Women's Preventive
Screening USPSTF Rating (Aug. 2021): B The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant persons at 24 weeks of gestation or after. For additional diabetes screening benefits, also see the Screening for Pre-Diabetes and Type 2 Diabetes row. Also see the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy.	Preventive Health section for Screening for Diabetes in Pregnancy codes.	Health section for Screening for Diabetes in Pregnancy preventive benefit instructions. Note: This benefit applies regardless of the gestational week.
Screening Mammography USPSTF Rating (2002): B The USPSTF recommends screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older. Also refer to the Medical Policy titled Breast Imaging for Screening and Diagnosing Cancer. Also see the Breast Cancer Screening for Average-Risk Women recommendation in the Expanded Women's Preventive Health section.	Procedure Code(s): 77063, 77067 Revenue Code: 0403 Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.	No age limits. Does not have diagnosis code requirements for the preventive benefit to apply. Note: This benefit only applies to screening mammography.

Preventive Care Services

United Healthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Cervical Cancer Screening

USPSTF Rating (Aug. 2018): A

Age 21-29 years:

The USPSTF recommends screening for cervical cancer every 3 years with cervical cytology alone in women aged 21-29 years.

Age 30-65 years:

For women aged **30 to 65 years**, the USPSTF recommends:

- Screening every 3 years with cervical cytology alone,
- Every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or
- Every 5 years with hrHPV testing in combination with cytology (co-testing).

Bright Futures, March 2014:

Adolescents should no longer be routinely screened for cervical dysplasia until age 21.

Also see <u>Screening for Cervical</u>
<u>Cancer</u> in the *Expanded Women's Preventive Health* section.

Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - Cholesterol Screening (Lipid Disorders Screening)

USPSTF Rating (August 2022): B

The USPSTF recommends that clinicians prescribe a statin for the primary prevention of CVD for adults aged 40 to 75 years who have 1 or more CVD risk factors (i.e. dyslipidemia, diabetes, hypertension, or smoking) and an estimated 10-year risk of a cardiovascular event of 10% or greater.

Code(s)

Human Papillomavirus DNA Testing (HPV)

Procedure Code(s):

0502U, 87624, 87625, 87626, G0476

Diagnosis Code(s):

Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4

Cervical Cytology (Pap Test) Code Group 1 Procedure Code(s):

G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, Q0091, P3000, P3001

Code Group 1 Diagnosis Code(s):

Does not have diagnosis code requirements for preventive benefit to apply.

Code Group 2 Procedure Code(s):

88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88155, 88164, 88165, 88166, 88167, 88174, 88175

Code Group 2 Diagnosis Code(s):

Z00.00, Z00.01, Z01.411, Z01.419, Z12.4

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Human Papillomavirus DNA Testing (HPV)

Age 30 years and up.

Requires one of the diagnosis codes listed in this row.

Cervical Cytology (Pap Test) Code Group 1:

Limited to age 21-65 years (ends on 66th birthday).

Does not have diagnosis code requirements for preventive benefits to apply.

Code Group 2:

Limited to age 21–65 years (ends on 66th birthday).

Requires one of the Code Group 2 diagnosis codes listed in this row.

Procedure Code(s):

Cholesterol Screening:

80061, 82465, 83718, 83719, 83721, 83722, 84478

Blood Draw:

36415, 36416

Risk Assessment: G0537

Diagnosis Code(s):

Z00.00, Z00.01, Z13.220

Cholesterol Screening:

Ages 40-75 years (ends on 76th birthday).

Requires one of the diagnosis codes listed in this row for 80061, 82465, 83718, 83719, 83721, 83722, 84478.

Blood Draw:

Ages 40-75 years (ends on 76th birthday): Requires one of the listed Cholesterol Screening procedure codes **and** one of the Diagnosis Codes listed in this row.

Risk Assessment:

Ages 40-75 years (ends on 76th birthday). The diagnosis codes listed in this row are not required for G0537

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Notes

- For statin medications benefits, refer to the pharmacy plan administrator.
- See <u>Dyslipidemia Screening</u> (<u>Bright Futures</u>) for recommendations for children.

Code(s)

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Preventive Benefit Does Not Apply:

For all ages above, if any of the following lipid disorders diagnosis codes are present in any position, the preventive benefit does **not** apply: E71.30, E75.5, E78.00, E78.01, E78.2, E78.3, E78.41, E78.49, E78.5, E78.79, E78.81, E78.89, E88.2, E88.89

Colorectal Cancer Screening

USPSTF Rating (May 2021): B

The USPSTF recommends screening for colorectal cancer in adults aged 45 to 49 years.

USPSTF Rating (May 2021): A

The USPSTF recommends screening for colorectal cancer in all adults aged 50 to 75 years.

Also refer to the Medical Policies titled <u>Outpatient Surgical</u>

<u>Procedures - Site of Service;</u>

<u>Screening Colonoscopy Procedures</u>

<u>- Site of Service;</u> and <u>Magnetic</u>

<u>Resonance Imaging (MRI) and</u>

<u>Computed Tomography (CT) Scan</u>

<u>- Site of Service.</u>

Also see the <u>Frequently Asked</u> Questions section.

Colonoscopy Procedure Code(s):

Preventive Colonoscopy: G0105. G0121

Preventive Colonoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right): 44388*, 44389*, 44392*, 44394*, 45378*, 45380*, 45381*, 45384*, 45385*, 45388*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Note: Also see the <u>Colonoscopy Pre-</u> Op Consultation row below.

Colonoscopy

Age Limits: 45-75 years (ends on 76th birthday).

Codes G0105 and G0121 do not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to G0104, G0105, G0121, G0328 or S0285

Sigmoidoscopy Procedure Code(s):

Preventive Sigmoidoscopy: G0104

Preventive Sigmoidoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right):

45330*, 45331*, 45333*, 45338*, 45346*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79,

Sigmoidoscopy

Age Limits: 45-75 years (ends on 76th birthday).

Code G0104 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to codes G0104, G0105, G0121, G0328 or S0285

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy)

Procedure Code(s):

Pathology: 88304, 88305

Anesthesia: 00811, 00812, 99152, 99153, 99156, 99157, G0500

Diagnosis Code(s):

Applies to the Pathology and Anesthesia codes listed above: Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Colonoscopy or Sigmoidoscopy)

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's

sections above for additional instructions.

Pathology and Anesthesia (for

Age Limits: 45-75 years (ends on 76th birthday).

Requires **both** of the following:

- One of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); and
- One of the procedure codes listed in the Colonoscopy row, or the Sigmoidoscopy row.

Note: Preventive benefits apply when the surgeon's claim is preventive.

Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT) **Procedure Code(s):**

Preventive:

G0328

Preventive When Billed with Certain Codes (see Preventive Benefit Instructions to the right): 82270*, 82274*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT)

Age Limits: 45-75 years (ends on 76th birthday).

Code G0328 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to G0104, G0105, G0121, G0328 or S0285.

Fecal DNA

Procedure Code(s):

0464U, 81528

Diagnosis Code(s):

Code 81528 does not have diagnosis code requirements for preventive benefits to apply.

Pre-Op Consultation Procedure Code(s):

Preventive: S0285

Fecal DNA

Age Limits: 45-75 years (ends on 76th birthday).

Benefit is limited to once every 3 years.

Codes 0464U and 81528 do not have diagnosis code requirements for preventive benefits to apply.

Pre-Op Consultation

Age Limits: 45-75 years (ends on 76th birthday).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Preventive when billed with one of the diagnosis codes listed in this row: 99202*, 99203*, 99204*, 99205*, 99211*, 99212*, 99213*, 99214*, 99215*, 99242*, 99243*, 99244*, 99245*, 99417*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above: Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Note: For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.

Preventive Benefit Instructions Refer to Coverage Rationale, and FA

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Code S0285 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when billed with one of the diagnosis codes listed in this row (Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79).

Computed Tomographic Colonography (Virtual Colonoscopy)

Procedure Code(s):

74263

Diagnosis Code(s):

Does not have diagnosis code requirements for preventive benefit to apply.

Computed Tomographic Colonography (Virtual Colonoscopy)

Age Limits: 45-75 years (ends on 76th birthday).

Does not have diagnosis code requirements for preventive benefit to apply.

Prior authorization requirements may apply, depending on plan.

Wellness Examinations

(well-baby, well-child, well-adult)

USPSTF Rating: None

UnitedHealthcare supports AAP and AAFP age and frequency guidelines.

HRSA Requirements:

The Wellness Examinations codes in this row include the following HRSA requirements for women, where applicable:

- Breastfeeding support, counseling, and education
- Contraceptive methods and sterilizations (counseling and follow-up care)

Procedure Code(s):

Medicare Wellness Exams: G0402, G0438, G0439

STIs Behavioral Counseling: G0445

Annual Gynecological Exams: S0610, S0612, S0613

Pelvic Examination (add-on code): 99459

Preventive Medicine Services (Evaluation and Management): 99381, 99382, 99383, 99384, 99385, 99386, 99387 99391, 99392, 99393, 99394, 99395, 99396, 99397 Does not have diagnosis code requirements for the preventive benefit to apply.

G0445 is limited to twice per year.

G0296 is limited to age 50 to 80 years (ends on 81st birthday).

Pelvic Examination add-on code 99459:

Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

- Screening and counseling for interpersonal domestic violence
- Screening for human immunodeficiency virus infection (HIV); education and risk assessment
- Counseling for sexually transmitted infections (STIs)
- Well-woman preventive visits
- Screening for urinary incontinence
- Obesity prevention in midlife women (counseling)

Code(s)

Preventive Medicine, Individual Counseling:

99401, 99402, 99403, 99404

Preventive Medicine, Group Counseling:

99411. 99412

Newborn Care (evaluation and management):

99461

Counseling Visit (to discuss the need for Lung Cancer Screening (LDCT) using Low Dose CT Scan): G0296

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Also see the Expanded Women's Preventive Health section.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Newborn Screenings

All newborns

USPSTF Rating (March 2008): A

Hypothyroidism Screening: Screening for congenital hypothyroidism in newborns.

USPSTF Rating (March 2008): A

Phenylketonuria Screening: Screening for phenylketonuria (PKU) in newborns.

USPSTF Rating (Sept. 2007): A

Sickle Cell Screening: Screening for sickle cell disease in newborns.

Note: For Bright Futures hearing screening, see Hearing Tests (Bright Futures).

Metabolic Screening Panel (Newborns)

Procedure Code(s):

Hypothyroidism Screening: 84437.84443

Phenylketonuria Screening: 84030, S3620

Sickle Cell Screening:

83020, 83021, 83030, 83033, 83051, S3850

Blood Draw:

36415, 36416

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Newborn Screenings:

Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.

Blood Draw:

Age 0-90 days, requires one of the listed Hypothyroidism Screening, Phenylketonuria Screening, or Sickle Cell Screening procedure codes.

Procedure Code(s):

Metabolic Screening Panel: 82017, 82136, 82261, 82775, 83020, 83498, 83516, 84030, 84437, 84443, S3620

Metabolic Screening Panel:

Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

For preventive care medications, refe	r to the pharmacy plan administrator.	
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
	Blood Draw: 36415, 36416 Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.	Blood Draw: Age 0-90 days. Requires one of the listed Metabolic Screening Panel procedure codes listed in this row.
USPSTF Rating (June 2018): B Women 65 and older: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 years and older. USPSTF Rating (June 2018): B Postmenopausal women younger than 65 years at increased risk of osteoporosis: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.	Procedure Code(s): 76977, 77080, 77081, G0130 Diagnosis Code(s): Z00.00, Z00.01, Z13.820, Z82.62	Requires one of the diagnosis codes listed in this row.
Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults USPSTF Rating (Nov. 2018): B The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.	Procedure Code(s): Alcohol or Drug Use Screening: 99408, 99409 Annual Alcohol Screening: G0442 Brief Counseling for Alcohol: G0443 Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.	Does not have diagnosis code requirements for preventive benefits to apply.

Also see the Expanded Women's Preventive Health section.

nes.

Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications, refer to the pharmacy plan administrator.					
	Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.		
	Bright Futures (April 2017): Bright Futures recommends alcohol or drug use assessments from age 11-21 years.				
	Also see rows: <u>Unhealthy Drug Use</u> <u>Screening (Adults)</u> ; and <u>Tobacco</u> , <u>Alcohol</u> , or <u>Drug Use Assessment</u> (Bright Futures).				
	Unhealthy Drug Use Screening (Adults) USPSTF Rating (June 2020): B	Procedure Code(s): Alcohol or Drug Use Screening: 99408, 99409	Does not have diagnosis code requirements for preventive benefits to apply.		
	The USPSTF recommends screening by asking questions	Diagnosis Code(s): Does not have diagnosis code			

screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.)

Bright Futures (April 2017):

Bright Futures recommends alcohol or drug use assessments from age 11-21 years.

Also see rows: Screening and **Behavioral Counseling Interventions** in Primary Care to Reduce Unhealthy Alcohol Use in Adults: and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures).

Does not have diagnosis code requirements for preventive benefit to apply.

High Blood Pressure in Adults – Screening

USPSTF Rating (April 2021): A

The USPSTF recommends screening for hypertension in adults 18 years or older with office blood pressure measurement.

The USPSTF recommends obtaining blood pressure measurements outside of the clinical setting for diagnostic

Blood Pressure Measurement in a Clinical Setting

N/A

Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting) **Procedure Code(s):**

Ambulatory Blood Pressure Measurement: 93784, 93786, 93788 or 93790

Blood Pressure Measurement in a Clinical Setting

This service is included in a preventive care wellness examination.

Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting)

Age 18 years and older. Requires the diagnosis code listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refe	r to the pharmacy plan administrator.	. ganao
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
confirmation before starting treatment.	Diagnosis Code(s): Abnormal Blood-Pressure Reading Without Diagnosis of Hypertension: R03.0	
Breast Cancer: Medication Use to Reduce Risk USPSTF Rating (Sept. 2019): B The USPSTF recommends that clinicians offer to prescribe risk- reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer and at low risk for adverse medication effects.	Procedure Code(s): Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, 99417, G0463 Diagnosis Code(s): Z80.3, Z80.41, Z15.01, Z15.02	Requires one of the diagnosis codes listed in this row in the primary position.
Primary Care Interventions to Promote Breastfeeding USPSTF Rating (Oct. 2016): B The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	N/A Also see the Expanded Women's Preventive Health section	Included in primary care or OB/GYN office visits.
Depression in Adults (Screening)	Procedure Code(s): 96127, 96161, G0136, G0444	Requires one of the diagnosis codes listed in this row, for 96127.
USPSTF Rating (June 2023): B The USPSTF recommends screening for depression in the adult population, including pregnant and postpartum persons, as well as older adults. Bright Futures (February 2017): Maternal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.	Diagnosis Code(s): Required for 96127 Only: Encounter for Screening for Depression: Z13.31, Z13.32	The diagnosis codes listed in this row are not required, for G0136, G0444, and 96161.
Also see the rows for <u>Screening for Anxiety</u> (HRSA); <u>Anxiety Disorders in Adults Screening</u> (USPSTF): <u>Depression in Children and Adolescents (Screening)</u> (USPSTF); <u>Perinatal Depression – Preventive Interventions</u>		

Preventive Care Services

Preventive Care Services Also see the Expanded Women's Preventive Health section. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications, refer to the pharmacy plan administrator. Service **Preventive Benefit Instructions** A date in this column is when the listed Refer to Coverage Rationale, and FAQ's rating was released, not when the Code(s) sections above for additional instructions. benefit is effective. (Counseling) (USPSTF): and Depression and Suicide Risk Screening (Bright Futures). Depression in Children Requires one of the diagnosis codes **Procedure Code(s):** listed in this row, for 96127. and Adolescents 96127, 96161, G0136, G0444 (Screening) The diagnosis codes listed in this row Diagnosis Code(s): are **not** required for G0136, G0444, and Required for 96127 Only: USPSTF Rating (October 2022): B 96161. Encounter for Screening for The USPSTF recommends Depression: screening for major depressive disorder (MDD) in adolescents aged Z13.31, Z13.32 12-18 years. **Bright Futures (February 2017):** Maternal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age. Note: The Bright Futures Periodicity Schedule recommends depression screening begin at age 12-21 years. Also see the rows for Anxiety Disorders in Adults Screening (USPSTF): Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures). Anxiety Disorders in Requires the diagnosis code listed in **Procedure Code(s):** this row. Adults (Screening) 96127 USPSTF Rating (June 2023): B **Diagnosis Code(s):** The USPSTF recommends Encounter for Screening Examination screening for anxiety in adults, for Other Mental Health and including pregnant and postpartum Behavioral Disorders: persons. This applies to adults age Z13.39 64 or younger. Also see the rows for Screening for Anxiety (HRSA); and Screening for Anxiety in Children and Adolescents (USPSTF).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Screening for Anxiety in Children and Adolescents

USPSTF Rating (October 2022): B The USPSTF recommends

screening for anxiety in children and adolescents aged 8 to 18 years.

Also see the rows for Anxiety Disorders in Adults Screening (USPSTF): Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal <u>Depression – Preventive</u> Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).

Code(s)

Procedure Code(s):

Diagnosis Code(s):

Encounter for Screening Examination for Other Mental Health and Behavioral Disorders:

Z13.39

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Requires the diagnosis code listed in this row.

Healthy Diet and Physical Activity for Cardiovascular Disease **Prevention in Adults** with Cardiovascular Risk Factors: Behavioral **Counseling Interventions**

USPSTF Rating (Nov. 2020): B

The USPSTF recommends offering or referring adults with cardiovascular disease risk factors to behavioral counseling interventions to promote a healthy diet and physical activity.

Procedure Code(s):

Medical Nutrition Therapy or Counseling:

97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886

Risk Assessment: G0537

Diagnosis Code(s):

Screening:

Z13.220

Nicotine Dependence, Tobacco Use, or Family History of IHD:

F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, Z87.891, Z82.49

Overweight:

E66.3, Z68.25, Z68.26, Z68.27, Z68.28. Z68.29

Body Mass Index 30.0 - 39.9: Z68.30, Z68.31, Z68.32, Z68.33,

Z68.34, Z68.35, Z68.36, Z68.37,

Z68.38, Z68.39

Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886

The diagnosis codes listed in this row are **not** required for G0446, G0447, G0473 and G0537.

G0446 is limited to once per year.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column is when the listed rating was released, not when the benefit is effective. Code(s) Body Mass Index 40.0 and Over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 Impaired Fasting Glucose: R73.01 Metabolic Syndrome; Insulin Resistance Syndrome Type A; Other Insulin Resistance: E88.810, E88.811, E88.818, E88.819	tionale, and FAQ's
Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 Impaired Fasting Glucose: R73.01 Metabolic Syndrome; Insulin Resistance Syndrome Type A; Other Insulin Resistance:	
R73.01 Metabolic Syndrome; Insulin Resistance Syndrome Type A; Other Insulin Resistance:	
Resistance Syndrome Type A; Other Insulin Resistance:	
Hyperlipidemia / Dyslipidemia: E78.00, E78.01, E78.1, E78.2, E78.3, E78.41, E78.49, E78.5	
Obesity: E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45	
Essential Hypertension:	
Resistant Hypertension: I1A.0 Secondary Hypertension: I15.0, I15.1, I15.2, I15.8, I15.9, N26.2	
Hypertension Complicating Pregnancy, Childbirth, and the Puerperium: O10.011, O10.012, O10.013, O10.019, O10.02, O10.03, O10.111, O10.112, O10.113, O10.119, O10.12, O10.13, O10.211, O10.212, O10.213, O10.219, O10.22, O10.23, O10.311, O10.312, O10.313, O10.319, O10.32, O10.33, O10.411, O10.412, O10.413, O10.419, O10.42, O10.43, O10.911, O10.912, O10.913, O10.919, O10.92, O10.93, O11.1, O11.2, O11.3, O11.4, O11.5, O11.9, O13.1, O13.2, O13.3, O13.4, O13.5, O13.9, O16.1, O16.2, O16.3, O16.4, O16.5, O16.9 Urgent/Emergency/Crisis Hypertension:	

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	m		

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Diabetes:

Diabetes Diagnosis Code List

Atherosclerosis:

Atherosclerosis Diagnosis Code List

Coronary Atherosclerosis:

125.10, 125.110, 125.111, 125.112, 125.118, 125.119, 125.700, 125.701, 125.702, 125.708, 125.709, 125.710, 125.720, 125.712, 125.718, 125.719, 125.720, 125.721, 125.722, 125.728, 125.729, 125.730, 125.731, 125.732, 125.738, 125.739, 125.750, 125.751, 125.752, 125.758, 125.759, 125.760, 125.761, 125.762, 125.762, 125.768, 125.769, 125.790, 125.791, 125.792, 125.798, 125.799, 125.810, 125.811, 125.812

Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral

USPSTF Rating (Sept. 2018): B

Interventions

The USPSTF recommends that clinicians offer or refer adults with a body mass index (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive multicomponent behavioral interventions.

Procedure Code(s):

Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886

Diagnosis Code(s):

Body Mass Index 30.0-39.9: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39

Body Mass Index 40.0 and over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Obesitv:

E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82

Procedure Code(s):

Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271

listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886

Requires one of the diagnosis codes

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's

sections above for additional instructions.

G0446 is limited to once per year. The diagnosis codes listed in this row are **not** required for G0446, G0447, and G0473.

High Body Mass Index in Children and Adolescents

USPSTF Rating (June 2024): B The USPSTF recommends that clinicians provide or refer children

Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886

G0446 is limited to once per year.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

÷	Δ	r۱	/1	^	Δ

A date in this column is when the listed rating was released, not when the benefit is effective.

and adolescents 6 years or older with a high body mass index (BMI) (≥ 95th percentile for age and sex) to comprehensive, intensive behavioral interventions. See the *Practice Considerations* section of the published USPSTF recommendation or more information about behavioral interventions.

Code(s)

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886

Also see the codes in the Wellness Examinations row above.

Diagnosis Code(s):

Obesity:

E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82

Pediatric BMI of 120% or more of the 95th percentile for age: Z68.54, Z68.55, Z68.56

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

The diagnosis codes listed in this row are **not** required for G0446, G0447, and G0473.

Healthy Weight and Weight Gain During Pregnancy: Behavioral Counseling Interventions

USPSTF Rating (May 2021): B

The USPSTF recommends that clinicians offer pregnant persons effective behavioral counseling interventions aimed at promoting healthy weight gain and preventing excess gestational weight gain in pregnancy.

Behavioral Counseling to Prevent Sexually Transmitted Infections

USPSTF Rating (Aug. 2020): B

The USPSTF recommends behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).

Procedure Code(s):

Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling: 99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: G0447, G0473

Diagnosis Code(s):

Pregnancy Diagnosis Codes

Procedure Code(s):

STIs Behavioral Counseling: G0445

Preventive Medicine Individual Counseling 99401, 99402, 99403, 99404

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Requires one of the diagnosis codes listed in this row.

Does not have diagnosis code requirements for the preventive benefit to apply.

G0445 is limited to twice per year.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Interventions for Tobacco Smoking Cessation in Adults, including Pregnant Persons

USPSTF Rating (Jan. 2021): A Pregnant Persons (A): The USPSTF recommends that clinicians ask all pregnant persons about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant persons who use tobacco.

Nonpregnant Adults (A): The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and US Food and Drug Administration (FDA)—approved pharmacotherapy for cessation to nonpregnant adults who use tobacco.

Note: Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.

Also see rows: <u>Unhealthy Drug Use</u>
<u>Screening (Adults)</u>; and <u>Tobacco</u>,
<u>Alcohol</u>, or <u>Drug Use Assessment</u>
(Bright Futures).

Primary Care Interventions To Prevent Tobacco Use In Children And Adolescents

USPSTF Rating (April 2013): B
The USPSTF recommends that
primary care clinicians provide
interventions, including education or
brief counseling, to prevent initiation
of tobacco use among school-aged
children and adolescents.

Bright Futures (April 2017):

Code(s)

Procedure Code(s):

Behavioral Interventions: 99406, 99407

Preventive Medicine, Individual Counseling: 99401, 99402, 99403, 99404

Also see the codes in the Wellness Examinations row above.

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Does not have diagnosis code requirements for the preventive benefit to apply.

Procedure Code(s):

Smoking and Tobacco Use Cessation Counseling Visit: 99406. 99407

Preventive Medicine, Individual Counseling: 99401, 99402, 99403, 99404
Also see the codes in the Wellness Examinations row above.

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Does not have diagnosis code requirements for the preventive benefit to apply.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Treventive care services			
Also see the Expanded Women's Preventive Health section.			
Certain codes may not be payable in all circumstances due to other policies or guidelines.			
For preventive care medications, refer to the pharmacy plan administrator.			
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
Bright Futures recommends tobacco use assessments from age 11-21 years.			
Also see rows: <u>Unhealthy Drug Use</u> <u>Screening (Adults)</u> ; and <u>Tobacco</u> , <u>Alcohol, or Drug Use Assessment</u> (<u>Bright Futures</u>).			
Screening for Visual Impairment in Children USPSTF Rating (Sept. 2017): B The USPSTF recommends vision screening at least once in all children age 3 to 5 years to detect amblyopia or its risk factors.	Procedure Code(s): Visual Acuity Screening (e.g., Snellen chart): 99173 Instrument-Based Screening: 99174, 99177	Visual Acuity Screening (99173): Up to age 21 years (ends on 22 nd birthday). Does not have diagnosis code requirements for preventive benefits to apply. Instrument-Based Screening (99174 and 99177):	
Bright Futures: Visual acuity screening is recommended for age 4 and 5 years as well as in cooperative 3 year olds. Instrument-based screening recommended for age 12 and 24 months, in addition to the well visits at 3-5 years of age.	Diagnosis Code(s): See the Preventive Benefit Instructions.	 Age 1 to 5 (ends on 6th birthday): Does not have diagnosis code requirements for preventive benefits to apply. Age 6 to 21 years (ends on 22nd birthday): Refer to the Medical Policy titled Ocular Photoscreening for allowable diagnoses. 	
Behavioral Counseling to Prevent Skin Cancer	N/A	This service is included in a preventive care wellness examination or focused E&M visit.	
USPSTF Rating (March 2018): B The USPSTF recommends counseling young adults, adolescents, children, and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons ages 6 months to 24 years with fair skin types to reduce their risk of skin cancer.			
Falls Prevention in Community-Dwelling Older Adults	N/A	This service is included in a preventive care wellness examination or focused E&M visit.	
USPSTF Rating (June 2024): B The USPSTF recommends exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.			

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Concerning for Intimate	
benefit is effective.	
rating was released, not when the	

A date in this column is when the listed

Code(s)

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Screening for Intimate Partner Violence

Service

USPSTF Rating (Oct. 2018): B
The USPSTF recommends that
clinicians screen for intimate partner
violence in women of reproductive
age and provide or refer women
who screen positive to ongoing
support services.

Also see <u>Screening and Counseling</u> <u>for Interpersonal and Domestic</u> <u>Violence</u> in the *Expanded Women's Preventive Health* section.

N/A

This service is included in a preventive care wellness examination.

Screening for Lung Cancer with Low-Dose Computed Tomography

USPSTF Rating (March 2021): B

The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 50 to 80 years who have a 20 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.

Procedure Code(s):

71271

Diagnosis Code(s):

F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891

Codes for Reporting Purposes:

G9275, G9276

Note: Codes G9275 and G9276, are for reporting purposes only, if applicable. These codes are not separately reimbursable.

Requires one of the diagnosis codes listed in this row.

Limitations:

- Limited to one per year, and
- All of the following criteria:
 - Age 50 to 80 years (ends on 81st birthday), and
 - At least 20 pack-years* of smoking history, and
 - Either a current smoker or has quit within the past 15 years

Note: Prior authorization requirements may apply, depending on plan.

*A pack-year is a way to measure the amount a person has smoked over a long period of time. It is calculated by multiplying the number of packs of cigarettes smoked per day by the number of years the person has smoked. For example, 1 pack year is equal to smoking 1 pack per day for 1 year, or 2 packs per day for half a year, and so on. Source: National Institutes of Health, National Cancer Institute
Dictionary of Cancer Terms, pack year definition web page.

https://www.cancer.gov/publications/dictionaries/cancer-terms/def/pack-year

Procedure Code(s):Application of Topical

Application of Topical Fluoride by Physician or Other Qualified Health Care Professional: 99188

Does not have diagnosis code requirements for the preventive benefit to apply.

Age 0-5years (ends on 6th birthday).

Fluoride Application in Primary Care

USPSTF Rating (May 2014): B

Preventive Care Services

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Children From Birth Through Age 5 Years. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.

Bright Futures (July 2022):

Bright Futures adopted the May 2014 recommendation of the USPSTF and further recommends. once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office, based on caries risk.

Code(s)

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Latent Tuberculosis Infection in Adults: Screening

USPSTF Rating (May 2023): B

The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk. This recommendation applies to asymptomatic adults 18 years or older at increased risk for tuberculosis (TB).

Procedure Code(s):

Screening: 86480, 86481, 86580

Follow-Up Visit to Check Results: 99211

Blood Draw: 36415, 36416

Diagnosis Code(s):

R76.11, R76.12, Z00.00, Z00.01, Z11.1, Z11.7, Z20.1

Note for age 18-21 years (ends on **22nd birthday)**: In addition to the codes in this row, the preventive benefit also applies to the diagnosis codes listed in the Bright Futures row: Tuberculosis (TB) Testing.

Screening:

Ages 18 years and up.

Requires one of the diagnosis codes listed in this row for CPT code 86480, 86481, and 86580.

Follow-Up Visit to Check Results (99211):

CPT code 99211 requires diagnosis code R76.11 or R76.12.

Blood Draw:

Ages 18 years and up.

Required to be billed with 86480 or 86481 and one of the diagnosis codes listed in this row.

Hypertensive Disorders of Pregnancy -Screening

USPSTF Rating (Sept. 2023): B

The USPSTF recommends screening for hypertensive disorders in pregnant persons with blood pressure measurements throughout pregnancy.

Hypertensive disorders of pregnancy screening by blood pressure measurement is included in the code for a prenatal care office visit. See the following code groups in the Expanded Women's Preventive Health section:

- **Prenatal Office Visits**
- **Prenatal Care Visits**
- **Global Obstetrical Codes**

See the following code groups in the Expanded Women's Preventive Health section:

- **Prenatal Office Visits**
- **Prenatal Care Visits**
- Global Obstetrical Codes

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Perinatal Depression – Preventive Interventions (Counseling)

USPSTF Rating (Feb. 2019): B

The USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal depression to counseling interventions.

Note: This policy addresses coding for interventions from a network medical provider only. For perinatal depression preventive interventions with a mental health provider, refer to the plan's mental health plan benefit administrator.

Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); and Depression Screening (Bright Futures).

Prevention of Human Immunodeficiency Virus (HIV) Infection: Preexposure Prophylaxis

USPSTF Rating (Aug. 2023): A

The USPSTF recommends that clinicians prescribe preexposure prophylaxis using effective antiretroviral therapy to persons who are at increased risk of HIV acquisition to decrease the risk of acquiring HIV.

Note: This benefit also includes:

- Kidney function testing (creatinine)
- Serologic testing for hepatitis B and C virus
- Testing for other STIs

Code(s)

Code Group 1 Procedure Code(s):

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Preventive Medicine, Group Counseling:

99411, 99412

Prenatal Care Visits: 59425, 59426

Preventive Medicine Services (Evaluation and Management): 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397

Code Group 2 Procedure Code(s):

Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0463

Code Group 2 Diagnosis Code(s):

A <u>Pregnancy Diagnosis Code</u>; **or** Z39.2 (encounter for routine postpartum follow-up); **or** Z13.32 (encounter for screening for maternal depression)

Procedure Code(s):

Kidney Function Testing (Creatinine): 82565, 82575

Pregnancy Testing: 81025, 84702, 84703

Office Visits:

99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463

(also see codes in the Wellness Examinations section)

Antiretroviral Therapy Injection: 96372 (Administration) J0739 (Injection cabotegravir, 1mg) G0012 (Administration)

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Code Group 1:

Does not have diagnosis code requirements for the preventive benefit to apply.

Code Group 2:

Requires one of the Code Group 2 diagnosis codes listed in this row.

Requires one of the diagnosis codes listed in this row in the primary position.

Note: Prior authorization requirements may apply, depending on plan. Refer to the Medical Benefit Drug Policy titled Long-Acting Injectable Antiretroviral Agents for HIV (for Commercial Only) or Long-Acting Injectable Antiretroviral Agents for HIV (for Individual Exchange Only).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

- Pregnancy testing when appropriate
- Ongoing follow-up and monitoring including HIV testing every 3 months

Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.

Code(s)

Counseling for PrEP to prevent HIV: G0011, G0013

Pharmacy Supplying Fee for HIV PrEP:

Q0521

Diagnosis Code(s):

Z11.3, Z11.4, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53
Also see the sections for:

- Behavioral Counseling to Prevent Sexually Transmitted Infections
- Chlamydia Infection Screening
- Gonorrhea Screening
- Hepatitis B Virus Infection Screening
- Hepatitis C Virus Infection Screening
- HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults
- Syphilis Screening

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Bright Futures

Anemia Screening in Children

(Bright Futures)

Procedure Code(s):

Anemia Screening in Children: 85014, 85018

Blood Draw: 36415, 36416

Diagnosis Code(s):

Z00.110, Z00.111, Z00.121, Z00.129, Z13.0

Anemia Screening in Children:

Ages prenatal to 21 (ends on 22nd birthday). No frequency limit.

Requires one of the diagnosis codes listed in this row.

Blood Draw:

Ages prenatal to 21 (ends on 22nd birthday).

Required to be billed with 85014 or 85018 **and** one of the diagnosis codes listed in this row.

Hearing Tests

Bright Futures (April 2017):

Hearing Tests: Recommended at ages: Newborn; between 3-5 days to 2 months; 4 years; 5 years, 6 years; 8 years; 10 years; once between age 11-14 years; once between age 15-17 years; once between age 18-21 years; also

Procedure Code(s):

Hearing Tests:

92551, 92552, 92553, 92558, 92587, 92588, 92650, 92651, V5008

Diagnosis Code(s):

Examination of Hearing: Z01.10 Routine Child: Z00.121, Z00.129 General Exam (for 18-21years): Z00.00, Z00.01 Ages 0-90 days: Does not have diagnosis code requirements for the preventive benefit to apply.

Ages 91 days to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row.
Limit of once per year.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

For preventive care medications, refer to the pharmacy plan administrator.			
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
recommended for those that have a positive risk assessment. Risk Assessment: Recommended at ages: 4 mo, 6 mo, 9 mo, 12 mo, 15 mo, 18 mo, 24 mo, 30 mo, 3 years, 7 years, and 9 years.	Note : A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.		
Screening for Visual Impairment in Children (Bright Futures)	See row above for <u>Screening for Visual Impairment in Children</u> .	See row above <u>Screening for Visual</u> <u>Impairment in Children</u> .	
Formal Developmental/ Autism Screening	Procedure Code(s): 96110	Ages prenatal to 2 years (ends on 3 rd birthday).	
 A formal, standardized developmental screen is recommended during the 9 month visit. A formal, standardized developmental screen is recommended during the 18 month visit, including a formal autism screen. A formal, standardized autism screen is recommended during the 24 month visit. A formal, standardized during the 24 month visit. A formal, standardized developmental screen is recommended during the 30 month visit. 	Diagnosis Code(s): Z00.121, Z00.129, Z13.40, Z13.41, Z13.42, Z13.49	No frequency limit. Requires one of the diagnosis codes listed in this row.	
Lead Screening Bright Futures: Screening Lab Work: Conduct risk assessment or screening, as appropriate, at the following intervals: 12 mo and 24 mo. Risk Assessment, and Screening if positive: Recommended at 6 mo, 9 mo, 12 mo, 18 mo, 24 mo, 3 years, 4 years, 5 years, and 6 years.	Procedure Code(s): Lead Screening: 83655 Blood Draw: 36415, 36416 Diagnosis Code(s): Z00.121, Z00.129, Z77.011	Lead Screening: Ages 6 months through age 6 years (ends on 7 th birthday). No frequency limit. Requires one of the diagnosis codes listed in this row. Blood Draw: Ages 6 months through age 6 years (ends on 7 th birthday). Required to be billed with 83655 and one of the diagnosis codes in this row.	
Tuberculosis (TB) Testing Bright Futures For age 18 years and older, also refer to the USPSTF recommendation above for Latent	Procedure Code(s): Screening: 86580 Follow-Up Visit to Check Results: 99211	Ages prenatal to 21 (ends on 22 nd birthday). Note: For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults No frequency limit.	

Preventive Care Services

United Healthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	m		

A date in this column is when the listed rating was released, not when the benefit is effective.

<u>Tuberculosis Infection: Screening,</u> Adults.

Code(s)

Diagnosis Code(s):

R76.11, R76.12, Z20.1, Z00.121, Z00.129, Z11.1, Z11.7

Note for age 18 years and older: In addition to these codes, the preventive benefit also applies to all codes listed in the USPSTF recommendation above for <u>Latent Tuberculosis Infection: Screening</u>, Adults.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

CPT code 86580 requires one of the diagnosis codes listed in this row.

CPT code 99211 requires diagnosis code R76.11, R76.12, or Z11.1.

Dyslipidemia Screening

Bright Futures (April 2014):

Risk Assessment: Recommended at 24 mo, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, 16 years.

Screening Lab Work: Conduct if risk assessment is positive, or, at the following intervals: once between age 9-11 years; once between age 17-21 years

Procedure Code(s):

Dyslipidemia Screening Lab Work: 80061, 82465, 83718, 83719, 83721, 83722, 84478

Blood Draw: 36415, 36416

Diagnosis Code(s):

Z00.121, Z00.129, Z13.220

Note: A risk assessment is included in the code for a wellness examination visit; see the Wellness Examinations row above.

Dyslipidemia Screening Lab Work:
Ages 24 months to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row.

Blood Draw:

Ages 24 months to 21 years (ends on 22nd birthday).

Requires one of the listed Dyslipidemia Screening procedure codes listed in this row **and** one of the diagnosis codes listed in this row.

Tobacco, Alcohol, or Drug Use Assessment

Bright Futures (April 2017):

Bright Futures recommends tobacco, alcohol, or drug use assessment from age 11-21 years.

See codes in the rows above:

- Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents
- Screening and Behavioral
 Counseling Interventions in
 Primary Care to Reduce
 Unhealthy Alcohol Use in Adults
- <u>Unhealthy Drug Use Screening</u> (Adults)

See the rows above:

- Primary Care Interventions To <u>Prevent Tobacco Use in Children</u> and Adolescents
- Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults
- Unhealthy Drug Use Screening (Adults)

Behavioral/Social/ Emotional Screening

Bright Futures (July 2022):

Bright Futures recommends behavioral/social/emotional screening annually from newborn to 21 years.

Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling)

An assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.

See the Wellness Examinations row above.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Page 32 of 46 Effective 04/01/2025

Preventive Care Services				
The state of the s	Also see the Expanded Women's Preventive Health section.			
Certain codes may not be payable in all circumstances due to other policies or guidelines.				
For preventive care medications, refe	r to the pharmacy plan administrator.			
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.		
(USPSTF); and <u>Depression and Suicide Risk Screening</u> (Bright Futures).				
Depression and Suicide Risk Screening	See the codes in the <u>Depression in</u> <u>Children and Adolescents</u> (Screening) row above.	See the <u>Depression in Children and</u> <u>Adolescents (Screening)</u> row above.		
Bright Futures (July 2022): Bright Futures recommends screening adolescents age 12-21 years for depression and suicide risk, making every effort to preserve confidentiality of the adolescent.				
Bright Futures (February 2017): Matemal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.				
Also see the rows for Screening for Anxiety (HRSA); Depression in Children and Adolescents (Screening) (USPSTF); and Perinatal Depression – Preventive Interventions (Counseling).				
Sexually Transmitted Infections (STI)	See the codes in the <u>Chlamydia</u> <u>Infection Screening</u> and <u>Gonorrhea</u> <u>Screening</u> rows above.	See the <u>Chlamydia Infection Screening</u> and <u>Gonorrhea Screening</u> rows above.		
Bright Futures (April 2017): Bright Futures recommends the following: STI Risk Assessment: Conduct risk assessment at each of the recommended visits between 11 years – 21 years. STI Lab Work: Conduct if risk assessment is positive.				
HIV Screening	See the codes in the HIV (Human	See the HIV (Human Immunodeficiency		
Bright Futures (April 2023): HIV Risk Assessment: Conduct risk assessment at age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years, and 21 years.	Immunodeficiency Virus) Screening for Adolescents and Adults row above.	Virus) Screening for Adolescents and Adults row above.		
HIV Screening Lab Work: Conduct at least once between age 15-21 years. Also recommended anytime between ages 11-14 years, when a				

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	r to the phannaey plan duminetrator.	
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.		
Sudden Cardiac Arrest (SCA) and Sudden Cardiac Death (SCD) – Risk Assessment and ECG Screening Bright Futures (July 2022): All children should be evaluated for conditions predisposing to SCA and SCD in the course of routine health care. A thorough and detailed history, family history, and physical examination are necessary to begin assessing SCA and SCD risk. The ECG should be the first test ordered when there is a concern for SCA risk. The ECG should be interpreted by a physician trained in recognizing electrical heart disease (i.e., a pediatric cardiologist or pediatric electrophysiologist).	Risk Procedure Code(s): 93000, 93005, 93010 Diagnosis Code(s): Required Screening Diagnosis Codes (requires at least one): Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 And requires one of the following Additional Diagnosis Codes (requires at least one): 142.0, 142.1, 142.2, 145.81, 149.8, 149.9, R55, R06.00, R06.09, R53.83, R00.2, R01.0, R01.1, R03.0, Q87.40, Q87.410, Q87.418, Q87.42, Q87.43, Q87.85, Q93.52, Z82.41, Z84.81, Z82.49 Risk Assessment A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.	ECG Screening for those at Risk: Limited to ages 11 years to 21 years (ends on 22 nd birthday). Requires one of the Screening Diagnosis Codes listed in this row and one of the Additional Diagnosis Codes listed in this row.
Hepatitis B Virus Infection Screening	See the codes in the <u>Hepatitis B</u> <u>Virus Infection Screening</u> row above.	See the <u>Hepatitis B Virus Infection</u> <u>Screening</u> row above.
Bright Futures (July 2022): Bright Futures recommends screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).		

Preventive Care Services Page 34 of 46 UnitedHealthcare Commercial and Individual Exchange Medical Policy

Proprietary Information of UnitedHealthcare. Copyright 2025 United HealthCare Services, Inc. Effective 04/01/2025

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Certain codes may not be payable in	all circumstances due to other policies o	r guidelines.
Service		Preventive Benefit Instructions
A date in this column reflects when the	2	Refer to Coverage Rationale, and FAQ's
listed rating was issued.	Code(s)	sections above for additional instructions.
Well-Woman Preventive	Procedure Code(s):	
Visits	Well-Woman Visits:	Well-Woman Visits:
		See the Wellness Examinations row in
HRSA Requirement (Dec. 2021): WPSI Recommends that women receive at least one preventive	See the Wellness Examinations row in the <i>Preventive Care Services</i> section.	the Preventive Care Services section.
care visit per year beginning in adolescence and continuing across the lifespan to ensure the provision of all recommended preventive services, including preconception and many services necessary for prenatal and interconception care, are obtained. The primary purpose of these visits should be the delivery and coordination of recommended preventive services as determined by age and risk factors. These services may be completed at a	Prenatal Office Visits: Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463	Prenatal Office Visits: Requires a Pregnancy Diagnosis Code.
	Pelvic Examination (add-on code): 99459	Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.
single or as part of a series of visits that take place over time to obtain all necessary services	Physician Prenatal Education, Group Setting: 99078	Physician Prenatal Education, Group Setting: Requires a Pregnancy Diagnosis Code.
depending on a woman's age, health status, reproductive health needs, pregnancy status, and risk factors. Well-women visits also include pre-pregnancy, prenatal, postpartum and interpregnancy visits. Also see Wellness Examinations and other USPSTF recommendations during pregnancy in the <i>Preventive Care Services</i> section.	Prenatal Care (Antepartum) Visits: 59425, 59426	Prenatal Care (Antepartum) Visits: Does not have diagnosis code requirements for the preventive benefit to apply.
	Global Obstetrical Codes: 59400, 59510, 59610, 59618	Global Obstetrical Codes: The routine, low-risk, prenatal visits portion of the code is covered as preventive. Does not have diagnosis code requirements for the preventive benefit to apply.
	Postpartum Care Visits (outpatient): 59430	Postpartum Care Visits (outpatient): Does not have diagnosis code requirements for the preventive benefit to apply.
	Diagnosis Code(s): Pregnancy Diagnosis Codes	See above services that require a pregnancy diagnosis code.
Screening for Diabetes in	Procedure Code(s):	Diabetes Screening:
Pregnancy	Diabetes Screening:	Requires a Pregnancy Diagnosis Code
HRSA Requirement (Jan. 2023):	82947, 82948, 82950, 82951, 82952, 83036	(regardless of gestational week).
Recommends screening pregnant		
women for gestational diabetes	Blood Draw:	
mellitus after 24 weeks of gestation (preferably between 24 and 28	36415, 36416	

Preventive Care Services

Page 35 of 46

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

, , ,
Service
A date in this column reflects when the listed rating was issued.
weeks of gestation) to prevent adverse birth outcomes. WPSI recommends screening pregnant women with risk factors for type 2 diabetes or GDM before 24 weeks of gestation – ideally at the first prenatal visit.
Also see the Screening for Pre-

<u>Diabetes and Type 2 Diabetes</u> and <u>Gestational Diabetes Screening</u> sections of the *Preventive Care Services* section, and the Screening for Diabetes After

Code(s)

Preventive Benefit InstructionsRefer to Coverage Rationale, and FAQ's

Diagnosis Code(s):

sections above for additional instructions.

Blood Draw:

Pregnancy Diagnosis Codes

Requires one of the diabetes screening procedure codes listed in this row **and** one of the Pregnancy Diagnosis Codes.

Note: If a diabetes diagnosis code is present in any position, the preventive benefit will **not** be applied. See the <u>Diabetes Diagnosis Code List</u>.

Pregnancy section. Screening for Diabetes After Pregnancy

HRSA Requirement (Jan. 2023):

Recommends screening for type 2 diabetes in women with a history of gestational diabetes mellitus (GDM) who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes. Initial testing should ideally occur within the first year postpartum and can be conducted as early as 4-6 weeks postpartum. Women who were not screened in the first year postpartum or those with a negative initial postpartum screening test result should be screened at least every 3 years for a minimum of 10 years after pregnancy. For those with a positive screening test result in the early postpartum period, testing should be repeated at least 6 months postpartum to confirm the diagnosis of diabetes regardless of the type of initial test (e.g., fasting plasma glucose, hemoglobin A1c, in the first 6 months postpartum regardless of whether the test results are positive or negative because the hemoglobin A1c test is less accurate during the first 6 months postpartum.

Procedure Code(s):

Diabetes Screening: 82947, 82948, 82950, 82951, 82952, 83036

Blood Draw: 36415, 36416

Diagnosis Code(s):

Required Screening Diagnosis Codes (requires at least one):

Z00.00, Z00.01, Z13.1

And requires the following additional code:

Additional Diagnosis Code Required: Z86.32 (personal history of gestational diabetes)

Diabetes Screening:

Requires one of the Required Screening diagnosis codes listed in this row **and** Z86.32.

No age limit.

Blood Draw:

Requires one of the Diabetes Screening procedure codes listed in this row **and** one of the Required Screening diagnosis codes listed in this row **and** 786.32.

Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the <u>Diabetes Diagnosis Code List</u>.

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above.

Certain codes may not be payable in all circumstances due to other policies or guidelines.		
Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
Also see Gestational Diabetes Screening and Screening for Pre- Diabetes and Type 2 in the Preventive Care Services section, and the Screening for Diabetes in Pregnancy section.		
Screening for Urinary Incontinence	See the Wellness Examinations row in the <i>Preventive Care Services</i>	See the Wellness Examinations row in the Preventive Care Services section
HRSA Requirement (Jan. 2024) The Women's Preventive Services Initiative recommends screening women for urinary incontinence annually. Screening should address whether women experience urinary incontinence and whether it impacts their activities and quality of life. If indicated, facilitating further evaluation and treatment is recommended.	section above.	above.
Counseling for Sexually Transmitted Infections (STIs)	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.
HRSA Requirement (Dec. 2021): WPSI recommends directed behavioral counseling by a health care clinician or other appropriately trained individual for sexually active adolescent and adult women at an increased risk for STIs. WPSI recommends that clinicians review a woman's sexual history and risk factors to help identify those at an increased risk of STIs. Risk factors include, but are not limited to, age younger than 25, a recent history of an STI, a new sex partner, multiple partners, a partner with concurrent partners, a partner with an STI, and a lack of or inconsistent condom use.		
For adolescents and women not identified as high risk, counseling to reduce the risk of STIs should be considered, as determined by clinical judgment.		

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service

A date in this column reflects when the listed rating was issued.

Screening for Human Immunodeficiency Virus Infection (HIV)

HRSA Requirement (Dec. 2021):

The Women's Preventive Services Initiative (WPSI) recommends all adolescent and adult women, ages 15 and older, receive a screening test for human immunodeficiency virus (HIV) at least once during their lifetime. Earlier or additional screening should be based on risk, and rescreening annually or more often may be appropriate beginning at age 13 for adolescent and adult women with an increased risk of HIV infection. The WPSI recommends risk assessment and prevention education for HIV infection beginning at age 13 and continuing as determined by risk. A screening test for HIV is recommended for all pregnant women upon initiation of prenatal care with rescreening during pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who present in labor with an undocumented HIV status.

Code(s)

Education and Risk Assessment

See the Wellness Examinations row in the *Preventive Care Services* section above

Screening Tests

See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the Preventive Care Services section above

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Education and Risk Assessment

See the <u>Wellness Examinations</u> row in the *Preventive Care Services* section above.

Screening Tests

See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the *Preventive Care Services* section above.

Contraceptive Methods (Including Sterilizations)

HRSA Requirement (Dec. 2021):

WPSI recommends that adolescent and adult women have access to the full range of contraceptives and contraceptive care to prevent unintended pregnancies and improve birth outcomes.

Contraceptive care includes screening, education, counseling, and provision of contraceptives (including in the postpartum period). Contraceptive care also includes follow-up care (e.g., management, evaluation, and changes, including the removal, continuation, and

Code Group 1 Procedure Code(s):

Sterilizations:

Tubal Ligation, Oviduct Occlusion:58600, 58605, 58611, 58615, 58670, 58671, A4264 (See Code Group 4 below for Tubal Ligation Follow-Up)

Contraceptive Methods:

Diaphragm or Cervical Cap: 57170, A4261, A4266
IUD (copper): J7300
IUD (Skyla®): J7301
IUD (Liletta®): J7297

IUD (Kyleena®): J7296 (See Code Group 2 below for

additional IUD codes)

Code Group 2 Procedure Code(s):

Code Group 1:

Does not have diagnosis code requirements for preventive benefits to apply.

Code Group 2:

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service

A date in this column reflects when the listed rating was issued.

discontinuation of contraceptives). WPSI recommends that the full range of U.S. Food and Drug Administration (FDA)-approved, granted, or -cleared contraceptives, effective family planning practices, and sterilization procedures be available as part of contraceptive care. The full range of contraceptives includes those currently listed in the FDA's Birth Control Guide: (1) sterilization surgery for women, (2) implantable rods, (3) copper intrauterine devices, (4) intrauterine devices with progestin (all durations and doses), (5) injectable contraceptives, (6) oral contraceptives (combined pill), (7) oral contraceptives (progestin only), (8) oral contraceptives (extended or continuous use), (9) the contraceptive patch, (10) vaginal contraceptive rings, (11) diaphragms, (12) contraceptive sponges, (13) cervical caps, (14) condoms. (15) spermicides. (16) emergency contraception (levonorgestrel), and (17) emergency contraception (ulipristal acetate), and any additional contraceptives approved, granted, or cleared by the FDA.

Notes:

- Coverage includes member reimbursement for the cost of FDA-approved, cleared, or granted mobile device applications for use as contraception consistent with the FDA-approved, cleared, or granted indication.
- For counseling and follow-up care, see the <u>Wellness</u>
 <u>Examinations</u> row in the *Preventive Care Services* section above.
- Certain employers may qualify for an exemption from covering contraceptive methods and

Code(s)

Contraceptive Methods:

Implantable Devices:

J7306, J7307

11976 (capsule removal)

11981 (implant insertion)

11982 (implant removal)

11983 (removal with reinsertion)

IUDs:

J7298 (Mirena®)

S4989

58300, S4981 (insertion)

58301 (removal)

(See Code Group 1 above for

additional IUD codes)

Injections:

96372 (administration)

J1050 (injection)

Code Group 2 Diagnosis Code(s):

These are required for Code Group 2. *Contraceptive Management:* Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9

Code Group 3 Procedure Code(s):

Anesthesia for Sterilization: 00851, 00940, 00942, 00950, 00952, 01960, 01961, 01965, 01966, 01967, 01968

Sterilization - Laparoscopy with Removal of Adnexal Structures: 58661

Code Group 3 Diagnosis Code(s):

This code is required for all Code Group 3 Procedure Codes: Sterilization:

Z30.2

Tubal Ligation Follow-Up Hysterosalpingogram Code Group 4 Procedure Code(s):

Catheterization and Introduction of Saline or Contrast Material: 58340

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Requires one of the Code Group 2 diagnosis codes listed in this row.

Code Group 3:

Requires one of the Code Group 3 diagnosis code listed in this row.

Code Group 4:

Requires one of the Code Group 4 diagnosis code listed in this row.

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above.

Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
sterilizations on account of religious objections. Refer to the plan's pharmacy benefit plan administrator for details on pharmacy contraceptives available under the plan's preventive benefit. Also refer to the Medical Policy titled Outpatient Surgical Procedures – Site of Service.	Hysterosalpingography: 74740 Contrast Material: Q9967 Code Group 4 Diagnosis Code(s): Tubal Ligation Status: Z98.51 Code Group 5 Procedure Code(s):	Code Group 5:	
	IUD Follow-Up Evaluation and Management (Office Visit): 99211, 99212 Pelvic Examination (add-on code): 99459 Refer to Code Group 7, Related Visits section below, for additional coding for Evaluation and Management (Office Visits). Code Group 5 Diagnosis Code(s): Encounter for routine checking of intrauterine contraceptive device: Z30.431	Requires one of the Code Group 5 diagnosis code listed in this row. Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.	
	Code Group 6 Procedure Code(s): Impacted IUD removal 58562 Code Group 6 Diagnosis Code(s): Z30.432, Z30.433	Code Group 6: Requires one of the Code Group 6 diagnosis codes listed in this row.	
	Code Group 7 Procedure Code(s): Related Visits: Related Evaluation and Management Office/Outpatient Visits for Contraception or Sterilization: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463 Pelvic Examination (add-on code): 99459 Also see coding in the Wellness Examinations row above.	Code Group 7: Requires one of the Code Group 7 diagnosis codes listed in this row. Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.	
	Related Pregnancy Tests:		

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or quidelines.

Certain codes may not be payable in	all circumstances due to other policies o	r guidelines.
Service		Preventive Benefit Instructions
A date in this column reflects when the	Codo(a)	Refer to Coverage Rationale, and FAQ's
listed rating was issued.	Code(s) Pregnancy Tests When Related to Contraception or Sterilization: 81025, 84702, 84703	sections above for additional instructions.
	Code Group 7 Diagnosis Codes: Tubal Ligation Status: Z98.51 Sterilization: Z30.2 Contraceptive Management: Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433,	
Breastfeeding Services	Z30.46, Z30.49, Z30.8, Z30.9 Counseling and Education	Counseling and Education
and Supplies	Procedure Code(s):	Requires one of the diagnosis codes
HRSA Requirement (Dec. 2021): WPSI recommends comprehensive lactation support services including consultation; counseling; education by clinicians and peer support services; and breastfeeding equipment and supplies) during the antenatal, perinatal, and postpartum periods to optimize the successful initiation and maintenance of breastfeeding. Breastfeeding equipment and supplies include, but are not limited to, double electric breast pumps (including pump parts and maintenance) and breast milk storage supplies. Access to double electric pumps should be a priority	98960, 98961, 98962, 99242*, 99243*, 99244*, 99245*, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, S9443 Also see the codes in the Wellness Examinations row in the Preventive Care Services section above. Diagnosis Code(s): B37.89, N61.1, N64.4, N64.51, N64.52, N64.53, N64.59, N64.89, O91.011, O91.012, O91.013, O91.019, O91.02, O91.03, O91.111, O91.112, O91.113, O91.211, O91.212, O91.213, O91.211, O91.212, O91.23, O92.011, O92.012, O92.013, O92.011, O92.012, O92.011, O92.012, O92.011, O92.013, O92.111, O92.112, O92.113,	listed in this row for 98960-98962, 99242-99245, 99341-99345, and 99347-99350. Does not have diagnosis code requirements for preventive benefits to apply for S9443.
to optimize breastfeeding and should not be predicated on prior failure of a manual pump. Breastfeeding equipment may also include equipment and supplies as clinically indicated to support dyads with breastfeeding difficulties and those who need additional services.	O92.119, O92.12, O92.13, O92.20, O92.29, O92.3, O92.4, O92.5, O92.70, O92.79, Q83.1, Q83.2, Q83.3, Q83.8,Z39.1, Z39.2 *For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.	
	Breastfeeding Equipment &	Breastfeeding Equipment &
	Supplies Procedure Code(s): Personal Use Electric Breast Pump: E0603	Supplies E0603 is limited to one purchase per birth. E0603 and A4281-A4287 require at
		least one of the diagnosis codes listed in

this row.

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Certain codes may not be payable in	all circumstances due to other policies of	r guidelines.
Service		Preventive Benefit Instructions
A date in this column reflects when the		Refer to Coverage Rationale, and FAQ's
listed rating was issued.	Code(s)	sections above for additional instructions.
	Breast Pump Supplies:	
	A4281, A4282, A4283, A4284,	
	A4285, A4286, A4287	
	Diagnosia Codo(a)	
	Diagnosis Code(s):	
	Pregnancy Diagnosis Codes or Z39.1.	
Screening and	See the Wellness Examinations row	See the Wellness Examinations row in
Counseling for	in the <i>Preventive Care Services</i> section above.	the <i>Preventive Care Services</i> section above.
Interpersonal and	section above.	above.
Domestic Violence		
HRSA Requirement (Dec. 2016):		
Recommends screening adolescents and women for		
interpersonal and domestic		
violence, at least annually, and,		
when needed, providing or referring		
for initial intervention services.		
Interpersonal and domestic		
violence includes physical violence,		
sexual violence, stalking and		
psychological aggression (including coercion), reproductive coercion,		
neglect, and the threat of violence,		
abuse, or both. Intervention		
services include, but are not limited		
to, counseling, education, harm		
reduction strategies, and referral to		
appropriate supportive services.		
Also see the Screening for Intimate		
Partner Violence row in the		
Preventive Care Services section		
above.		
Breast Cancer Screening	See the Screening Mammography	See the Screening Mammography row
for Average-Risk Women	row in the Preventive Care Services	in the Preventive Care Services section
	section above.	above.
HRSA Requirement (Dec. 2016):		
Recommends that average-risk		
women initiate mammography		
screening no earlier than age 40		
and no later than age 50.		
Screening mammography should occur at least biennially and as		
frequently as annually. Screening		
should continue through at least		
age 74 and age alone should not		
be the basis to discontinue		
screening. These screening		
recommendations are for women at		

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or quidelines.

Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
average risk of breast cancer. Women at increased risk should also undergo periodic mammography screening; however, recommendations for additional services are beyond the scope of this recommendation.			
Screening for Cervical Cancer	Human Papillomavirus DNA Testing (HPV)	Human Papillomavirus DNA Testing (HPV)	
HRSA Requirement (Dec. 2016): Recommends cervical cancer	See the <u>Cervical Cancer Screening</u> row in the <i>Preventive Care Services</i> section above.	See the <u>Cervical Cancer Screening</u> row in the <i>Preventive Care Services</i> section above.	
screening for average-risk women aged 21 to 65 years. For women aged 21 to 29 years recommends cervical cancer screening using cervical cytology (Pap test) every 3 years. Co-testing with cytology and human papillomavirus testing is not recommended for women younger than 30 years. Women aged 30 to 65 years should be screened with cytology and human papillomavirus testing every 5 years or cytology alone every 3 years. Women who are at average risk should not be screened more than once every 3 years.	Cervical Cytology (Pap Test) See the Cervical Cancer Screening row in the Preventive Care Services section above.	Cervical Cytology (Pap Test) See the Cervical Cancer Screening row in the Preventive Care Services section above.	
Screening for Anxiety HRSA Requirement (Dec. 2019): The Women's Preventive Services Initiative recommends screening for anxiety in adolescent and adult women, including those who are pregnant or postpartum. Optimal screening intervals are unknown and clinical judgement should be used to determine screening frequency. Given the high prevalence of anxiety disorders, lack of recognition in clinical practices, and multiple problems associated with untreated anxiety, clinicians should consider screening women who have not been recently screened. Also see the rows for Anxiety Disorders in Adults Screening (USPSTF); Screening for Anxiety in	Procedure Code(s): 96127 Diagnosis Code(s): Encounter for Screening Examination for Other Mental Health and Behavioral Disorders: Z13.39	Requires the diagnosis code listed in this row.	

Preventive Care Services

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the Preventive Care Services section above.

Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
Children and Adolescents (USPSTF); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression Screening (Bright Futures) in the Preventive Care Services section above.			
Obesity Prevention in Midlife Women (Counseling)	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.	
HRSA Requirement (Dec. 2021): WPSI recommends counseling midlife women aged 40 to 60 years with normal or overweight body mass index (BMI) (18.5-29.9 km/m2) to maintain weight or limit weight gain to prevent obesity. Counseling may include individualized discussion of healthy eating and physical activity.			

Diagnosis Codes

Preventive Care Services: ICD-10 Diagnosis Codes

Preventive Vaccine Codes

Preventive Care Services: Vaccine Codes

Benefit Considerations

Certain plans are not required to include coverage for the services identified by the federal Patient Protection and Affordable Care Act (PPACA). Refer to the member specific benefit plan document for coverage details.

Clinical Evidence

Refer to the Service column in the Applicable Codes section for the recommendation statements supporting this policy.

References

ACIP Vaccine-Specific Recommendations: https://www.cdc.gov/acip-recs/hcp/vaccinespecific/?CDC AAref Val=https://www.cdc.gov/vaccines/hcp/acip-recs/index.html. Accessed January 3, 2025.

American Academy of Family Physicians (AAFP) Clinical Preventive Services Recommendations: https://www.aafp.org/family-physician/patient-care/clinical-recommendations/clinical-practice-guidelines/clinicalpreventive-services-recommendations.html. Accessed January 3, 2025.

American Academy of Pediatrics, Bright Futures Guidelines, 4th edition, Evidence and Rationale chapter: https://www.aap.org/en/practice-management/bright-futures/bright-futures-materials-and-tools/bright-futures-guidelines-and-pocket-guide/. Accessed January 3, 2025.

American Academy of Pediatrics/Bright Futures/Recommendations for Pediatric Preventive Healthcare. (For ages 0-21): https://downloads.aap.org/AAP/PDF/periodicity_schedule.pdf. Accessed January 3, 2025.

American Academy of Pediatrics: http://www.aap.org/. Accessed January 3, 2025.

Centers for Disease Control and Prevention/Immunization Schedules: https://www.cdc.gov/vaccines/hcp/imz-schedules/index.html. Accessed January 3, 2025.

Grade Definitions for USPSTF Recommendations: http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions. Accessed January 3, 2025.

July 19, 2010 IRS Interim Rules: http://www.irs.gov/irb/2010-29 IRB/index.html. Accessed January 3, 2025.

Published Recommendations, U.S. Preventive Services Task Force:

http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations. Accessed January 3, 2025.

U.S. Food and Drug Administration (FDA), Vaccines Licensed for Use in the United States: http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM093833. Accessed January 3, 2025.

U.S. Vaccine Names Information Page: https://www.cdc.gov/vaccines/hcp/vaccines-us/index.html. Accessed January 3, 2025.

Women's Preventive Services Guidelines (HRSA): https://www.hrsa.gov/womens-guidelines. Accessed January 3, 2025.

Women's Preventive Services Initiative (WPSI): https://www.womenspreventivehealth.org/recommendations/. Accessed January 3, 2025.

Policy History/Revision Information

Dete	Oursease of Oherses	
2000	· · · · · · · · · · · · · · · · · · ·	
Date 04/01/2025	Applicable Codes Added instruction to refer to the Coverage Rationale and Frequently Asked Questions (FAQ) sections of the policy for additional preventive benefit instructions Preventive Care Services Screening for Pre-Diabetes and Type 2 Diabetes; Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors: Behavioral Counseling Interventions; Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral Interventions; High Body Mass Index in Children and Adolescents; and Healthy Weight and Weight Gain During Pregnancy: Behavioral Counseling Interventions Removed HCPCS code S9470 Colorectal Cancer Screening	
	 Removed coverage guidelines for Barium Enema (HCPCS codes G0106, G0120, and G0122) Colonoscopy; Sigmoidoscopy; and Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT) Updated preventive benefit instructions; removed language indicating colonoscopy screening procedure codes are preventive when billed in addition to HCPCS codes G0106, G0120, and G0122 Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy) Updated list of applicable CPT codes for Anesthesia; added 00811 Supporting Information Archived previous policy version MP.016.52 	

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates.

UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence (Medicare IOM Pub. No. 100-16, Ch. 4, §90.5).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.