

Spinal Fusion and Decompression (for Idaho Only)

Policy Number: CS365ID.A
Effective Date: June 1, 2025

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Medical Records Documentation Used for Reviews	2
Definitions	2
Applicable Codes	3
Description of Services	9
Clinical Evidence	9
U.S. Food and Drug Administration	11
References	11
Policy History/Revision Information	12
Instructions for Use	12

Related Policies
• Discogenic Pain Treatment (for Idaho Only)
• Epidural Steroid Injections for Spinal Pain (for Idaho Only)
• Facet Joint and Medial Branch Block Injections for Spinal Pain (for Idaho Only)
• Spinal Fusion and Bone Healing Enhancement Products (for Idaho Only)
• Total Artificial Disc Replacement for the Spine (for Idaho Only)

Application

This Medical Policy only applies the state of Idaho, including Idaho Medicaid Plus plans.

Coverage Rationale

Spinal procedures for the treatment of spine pain are proven and medically necessary in certain circumstances.

For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures:

- Decompression +/- Fusion, Cervical
- Decompression +/- Fusion, Lumbar
- Decompression +/- Fusion, Thoracic
- Fusion, Cervical Spine
- Fusion, Lumbar Spine
- Fusion, Thoracic Spine
- Scoliosis or Kyphosis Surgery
- Scoliosis or Kyphosis Surgery (Pediatric)

[Click here to view the InterQual® criteria.](#)

Isolated Facet Fusion, with or without instrumentation, is unproven and not medically necessary due to insufficient evidence of efficacy.

Dynamic Stabilization systems for the treatment of degenerative **Spondylolisthesis** are unproven and not medically necessary due to insufficient evidence of efficacy.

Total Facet Joint Arthroplasty is unproven and not medically necessary due to insufficient evidence of efficacy.

Dividing treatment of symptomatic, multi-site spinal pathology via anterior or posterior approach into serial or **Staged Multiple Sessions** when one session can address all sites is unproven and not medically necessary due to insufficient evidence of safety and efficacy.

Medical Records Documentation Used for Reviews

Benefit coverage for health services is determined by the federal, state, or contractual requirements, and applicable laws that may require coverage for a specific service. Medical records documentation may be required to assess whether the member meets the clinical criteria for coverage but does not guarantee coverage of the services requested; refer to the guidelines titled [Medical Records Documentation Used for Reviews](#).

Definitions

Disabling Symptoms: Are defined in a pivotal study demonstrating benefit of surgery (Weinstein, 2009), where the participants with an Oswestry Disability Index score of more than 8, or an SF-36 Bodily Pain Score of less than 70 or a Physical Function Score of less than 78 were the ones that demonstrated benefit. These scores are equal to or more severe than the majority of participants, meaning those participants within two standard deviations (+/-) of the mean for such scores.

Dynamic Stabilization: Also known as soft stabilization or flexible stabilization has been proposed as an adjunct or alternative to Spinal Fusion for the treatment of severe refractory pain due to degenerative Spondylolisthesis, or continued severe refractory back pain following prior fusion, sometimes referred to as failed back surgery syndrome. Dynamic Stabilization uses flexible materials rather than rigid devices to stabilize the affected spinal segment(s). These flexible materials may be anchored to the vertebrae by synthetic cords or by pedicle screws. Unlike the rigid fixation of Spinal Fusion, Dynamic Stabilization is intended to preserve the mobility of the spinal segment (Veritas Health, 2022).

Isolated Facet Fusion: A minimally invasive back procedure that uses specially designed bone dowels made from allograft material (donated cortical bone) that are inserted into the facet joints. The procedure is designed to stop facet joints from moving and is intended to eliminate or reduce back pain caused by facet joint dysfunction (Gellhorn, 2013).

Lumbar Spinal Stenosis (LSS): Narrowing or constriction of the lumbar spinal canal that may result in painful compression of a nerve and/or blood vessel(s) supplying the nerve (Veritas Health, 2022).

Progressive: Significant worsening of deficits or symptoms based on at least two measurements over days or weeks (rapidly Progressive) or over months (Progressive) on a validated pain or function scale or quantifiable symptoms (Veritas Health, 2022).

Radicular Pain: Pain which radiates from the spine into the extremity along the course of the spinal nerve root. The pain should follow the pattern of a dermatome associated with the irritated nerve root identified (Lenahan, 2018).

- Presenting symptoms should include a positive nerve root tension sign (positive straight leg raise test or femoral tension sign), or a reflex (asymmetric depressed reflex), sensory (asymmetric decreased sensation in a dermatomal distribution), or motor (asymmetric weakness in a myotomal distribution) deficit that correspond to the specific affected nerve root (Birkmeyer, 2002).
- As surgery is meant to relieve Radicular Pain from nerve root compression, imaging should show compression of the corresponding nerve root. (Lenahan, 2018).

Spinal Fusion: Also called Arthrodesis, is a surgical technique that may be done as an open or minimally invasive procedure. There are many different approaches to Spinal Fusion, but all techniques involve removing the disc between two or more vertebrae and fusing the adjacent vertebrae together using bone grafts and/or spacers placed where the disc used to be. Spacers can be made of bone or bone substitutes, metal (titanium), carbon fiber, polymers or bioresorbable materials and are often supported by plates, screws, rods, and/or cages (Veritas Health, 2022).

Spondylolisthesis: An acquired condition that involves the anterior displacement of one vertebral segment over subjacent vertebrae. The causes can be congenital, due to stress fractures, facet degeneration, injury, or after decompression surgery. The condition may be asymptomatic, or cause significant pain and nerve-related symptoms. If the slippage occurs backwards, it is referred to as retrolisthesis and lateral slippage is called listhesis (NASS, 2014a). Listhesis demonstrated on imaging is considered clinically significant (as opposed to a normal age-related change without clinical implication) if sagittal plane displacement is at least 3 mm on flexion and extension views or relative sagittal plane angulation greater than 11 degrees (Ghogawala et al, 2016).

Spondylolysis: A bone defect in the pars interarticularis; the isthmus or bone bridges between the inferior and superior articular surfaces of the neural arch of single vertebrae, most often the result of a stress fracture nonunion. The condition is an acquired condition, occurs commonly at a young age and may occur with or without Spondylolisthesis. The main

presenting symptom is back pain which in children is often treated with conservatively with orthotic bracing, activity modification and physical therapy. In adults' treatment involves education, analgesics and NSAIDS, with exercise and rapid return to activities. Once Spondylolisthesis occurs healing of the pars is unlikely. Surgery is indicated when there is Progressive neurological deficit, cauda equina compression, or persistent severe leg and back pain despite aggressive conservative management (Spinelli, 2008).

Staged Multi Session: Includes procedures performed on different days or requiring an additional anesthesia session.

Total Facet Arthroplasty: The implantation of a spinal prosthesis to restore posterior element structure and function, as an adjunct to neural decompression (Veritas Health, 2022).

Unremitting: Constant and unrelieved by conservative therapy (Veritas Health, 2022).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
0202T	Posterior vertebral joint(s) arthroplasty (e.g., facet joint[s] replacement) including facetectomy, laminectomy, foraminotomy, and vertebral column fixation, injection of bone cement, when performed including fluoroscopy, single level, lumbar spine
0219T	Placement of a posterior intrafacet implant(s), unilateral or bilateral, including imaging and placement of bone graft(s) or synthetic device(s), single level; cervical
0220T	Placement of a posterior intrafacet implant(s), unilateral or bilateral, including imaging and placement of bone graft(s) or synthetic device(s), single level; thoracic
0221T	Placement of a posterior intrafacet implant(s), unilateral or bilateral, including imaging and placement of bone graft(s) or synthetic device(s), single level; lumbar
0222T	Placement of a posterior intrafacet implant(s), unilateral or bilateral, including imaging and placement of bone graft(s) or synthetic device(s), single level; each additional vertebral segment (List separately in addition to code for primary procedure)
0719T	Posterior vertebral joint replacement, including bilateral facetectomy, laminectomy, and radical discectomy, including imaging guidance, lumbar spine, single segment
22206	Osteotomy of spine, posterior or posterolateral approach, 3 columns, 1 vertebral segment (e.g., pedicle/vertebral body subtraction); thoracic
22207	Osteotomy of spine, posterior or posterolateral approach, 3 columns, 1 vertebral segment (e.g., pedicle/vertebral body subtraction); lumbar
22208	Osteotomy of spine, posterior or posterolateral approach, 3 columns, 1 vertebral segment (e.g., pedicle/vertebral body subtraction); each additional vertebral segment (List separately in addition to code for primary procedure)
22210	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; cervical
22212	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; thoracic
22214	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; lumbar
22216	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; each additional vertebral segment (List separately in addition to primary procedure)
22220	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment; cervical
22222	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment; thoracic
22224	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment; lumbar
22226	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment; each additional vertebral segment (List separately in addition to code for primary procedure)

CPT Code	Description
22532	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic
22533	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); lumbar
22534	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic or lumbar, each additional vertebral segment (List separately in addition to code for primary procedure)
22548	Arthrodesis, anterior transoral or extraoral technique, clivus-C1-C2 (atlas-axis), with or without excision of odontoid process
22551	Arthrodesis, anterior interbody, including disc space preparation, discectomy, osteophylectomy and decompression of spinal cord and/or nerve roots; cervical below C2
22552	Arthrodesis, anterior interbody, including disc space preparation, discectomy, osteophylectomy and decompression of spinal cord and/or nerve roots; cervical below C2, each additional interspace (List separately in addition to code for separate procedure)
22554	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); cervical below C2
22556	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic
22558	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); lumbar
22585	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); each additional interspace (List separately in addition to code for primary procedure)
22590	Arthrodesis, posterior technique, craniocervical (occiput-C2)
22595	Arthrodesis, posterior technique, atlas-axis (C1-C2)
22600	Arthrodesis, posterior or posterolateral technique, single interspace; cervical below C2 segment
22610	Arthrodesis, posterior or posterolateral technique, single interspace; thoracic (with lateral transverse technique, when performed)
22612	Arthrodesis, posterior or posterolateral technique, single interspace; lumbar (with lateral transverse technique, when performed)
22614	Arthrodesis, posterior or posterolateral technique, single interspace; each additional interspace (List separately in addition to code for primary procedure)
22630	Arthrodesis, posterior interbody technique, including laminectomy and/or discectomy to prepare interspace (other than for decompression), single interspace, lumbar
22632	Arthrodesis, posterior interbody technique, including laminectomy and/or discectomy to prepare interspace (other than for decompression), single interspace, lumbar; each additional interspace (List separately in addition to code for primary procedure)
22633	Arthrodesis, combined posterior or posterolateral technique with posterior interbody technique including laminectomy and/or discectomy sufficient to prepare interspace (other than for decompression), single interspace, lumbar
22634	Arthrodesis, combined posterior or posterolateral technique with posterior interbody technique including laminectomy and/or discectomy sufficient to prepare interspace (other than for decompression), single interspace, lumbar; each additional interspace (List separately in addition to code for primary procedure)
22800	Arthrodesis, posterior, for spinal deformity, with or without cast; up to 6 vertebral segments
22802	Arthrodesis, posterior, for spinal deformity, with or without cast; 7 to 12 vertebral segments
22804	Arthrodesis, posterior, for spinal deformity, with or without cast; 13 or more vertebral segments
22808	Arthrodesis, anterior, for spinal deformity, with or without cast; 2 to 3 vertebral segments
22810	Arthrodesis, anterior, for spinal deformity, with or without cast; 4 to 7 vertebral segments
22812	Arthrodesis, anterior, for spinal deformity, with or without cast; 8 or more vertebral segments
22830	Exploration of spinal fusion

CPT Code	Description
22840	Posterior non-segmental instrumentation (e.g., Harrington rod technique, pedicle fixation across 1 interspace, atlantoaxial transarticular screw fixation, sublaminar wiring at C1, facet screw fixation) (List separately in addition to code for primary procedure)
22841	Internal spinal fixation by wiring of spinous processes (List separately in addition to code for primary procedure)
22842	Posterior segmental instrumentation (e.g., pedicle fixation, dual rods with multiple hooks and sublaminar wires); 3 to 6 vertebral segments (List separately in addition to code for primary procedure)
22843	Posterior segmental instrumentation (e.g., pedicle fixation, dual rods with multiple hooks and sublaminar wires); 7 to 12 vertebral segments (List separately in addition to code for primary procedure)
22844	Posterior segmental instrumentation (e.g., pedicle fixation, dual rods with multiple hooks and sublaminar wires); 13 or more vertebral segments (List separately in addition to code for primary procedure)
22845	Anterior instrumentation; 2 to 3 vertebral segments (List separately in addition to code for primary procedure)
22846	Anterior instrumentation; 4 to 7 vertebral segments (List separately in addition to code for primary procedure)
22847	Anterior instrumentation; 8 or more vertebral segments (List separately in addition to code for primary procedure)
22848	Pelvic fixation (attachment of caudal end of instrumentation to pelvic bony structures) other than sacrum (List separately in addition to code for primary procedure)
22849	Reinsertion of spinal fixation device
22850	Removal of posterior nonsegmental instrumentation (e.g., Harrington rod)
22852	Removal of posterior segmental instrumentation
22853	Insertion of interbody biomechanical device(s) (e.g., synthetic cage, mesh) with integral anterior instrumentation for device anchoring (e.g., screws, flanges), when performed, to intervertebral disc space in conjunction with interbody arthrodesis, each interspace (List separately in addition to code for primary procedure)
22854	Insertion of intervertebral biomechanical device(s) (e.g., synthetic cage, mesh) with integral anterior instrumentation for device anchoring (e.g., screws, flanges), when performed, to vertebral corpectomy(ies) (vertebral body resection, partial or complete) defect, in conjunction with interbody arthrodesis, each contiguous defect (List separately in addition to code for primary procedure)
22855	Removal of anterior instrumentation
22859	Insertion of intervertebral biomechanical device(s) (e.g., synthetic cage, mesh, methylmethacrylate) to intervertebral disc space or vertebral body defect without interbody arthrodesis, each contiguous defect (List separately in addition to code for primary procedure)
22899	Unlisted procedure, spine
62380	Endoscopic decompression of spinal cord, nerve root(s), including laminotomy, partial facetectomy, foraminotomy, discectomy and/or excision of herniated intervertebral disc, 1 interspace, lumbar
63001	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), 1 or 2 vertebral segments; cervical
63003	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), 1 or 2 vertebral segments; thoracic
63005	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), 1 or 2 vertebral segments; lumbar, except for spondylolisthesis
63012	Laminectomy with removal of abnormal facets and/or pars inter-articularis with decompression of cauda equina and nerve roots for spondylolisthesis, lumbar (Gill type procedure)
63015	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), more than 2 vertebral segments; cervical

CPT Code	Description
63016	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), more than 2 vertebral segments; thoracic
63017	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (e.g., spinal stenosis), more than 2 vertebral segments; lumbar
63020	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc; 1 interspace, cervical
63030	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc; 1 interspace, lumbar
63035	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc; each additional interspace, cervical or lumbar (List separately in addition to code for primary procedure)
63040	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; cervical
63042	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; lumbar
63043	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; each additional cervical interspace (List separately in addition to code for primary procedure)
63044	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; each additional lumbar interspace (List separately in addition to code for primary procedure)
63045	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [e.g., spinal or lateral recess stenosis]), single vertebral segment; cervical
63046	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [e.g., spinal or lateral recess stenosis]), single vertebral segment; thoracic
63047	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [e.g., spinal or lateral recess stenosis]), single vertebral segment; lumbar
63048	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [e.g., spinal or lateral recess stenosis]), single vertebral segment; each additional vertebral segment, cervical, thoracic, or lumbar (List separately in addition to code for primary procedure)
63050	Laminoplasty, cervical, with decompression of the spinal cord, 2 or more vertebral segments;
63051	Laminoplasty, cervical, with decompression of the spinal cord, 2 or more vertebral segments; with reconstruction of the posterior bony elements (including the application of bridging bone graft and non-segmental fixation devices [e.g., wire, suture, mini-plates], when performed)
63052	Laminectomy, facetectomy, or foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s] [e.g., spinal or lateral recess stenosis]), during posterior interbody arthrodesis, lumbar; single vertebral segment (List separately in addition to code for primary procedure)
63053	Laminectomy, facetectomy, or foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s] [e.g., spinal or lateral recess stenosis]), during posterior interbody arthrodesis, lumbar; each additional vertebral segment (List separately in addition to code for primary procedure)
63055	Transpedicular approach with decompression of spinal cord, equina and/or nerve root(s) (e.g., herniated intervertebral disc), single segment; thoracic

CPT Code	Description
63056	Transpedicular approach with decompression of spinal cord, equina and/or nerve root(s) (e.g., herniated intervertebral disc), single segment; lumbar (including transfacet, or lateral extraforaminal approach) (e.g., far lateral herniated intervertebral disc)
63057	Transpedicular approach with decompression of spinal cord, equina and/or nerve root(s) (e.g., herniated intervertebral disc), single segment; each additional segment, thoracic or lumbar (List separately in addition to code for primary procedure)
63064	Costovertebral approach with decompression of spinal cord or nerve root(s), (e.g., herniated intervertebral disc), thoracic; single segment
63066	Costovertebral approach with decompression of spinal cord or nerve root(s), (e.g., herniated intervertebral disc), thoracic; each additional segment (List separately in addition to code for primary procedure)
63075	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophyctomy; cervical, single interspace
63076	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophyctomy; cervical, each additional interspace (List separately in addition to code for primary procedure)
63077	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophyctomy; thoracic, single interspace
63078	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophyctomy; thoracic, each additional interspace (List separately in addition to code for primary procedure)
63081	Vertebral corpectomy (vertebral body resection), partial or complete, anterior approach with decompression of spinal cord and/or nerve root(s); cervical, single segment
63082	Vertebral corpectomy (vertebral body resection), partial or complete, anterior approach with decompression of spinal cord and/or nerve root(s); cervical, each additional segment (List separately in addition to code for primary procedure)
63085	Vertebral corpectomy (vertebral body resection), partial or complete, transthoracic approach with decompression of spinal cord and/or nerve root(s); thoracic, single segment
63086	Vertebral corpectomy (vertebral body resection), partial or complete, transthoracic approach with decompression of spinal cord and/or nerve root(s); thoracic, each additional segment (List separately in addition to code for primary procedure)
63087	Vertebral corpectomy (vertebral body resection), partial or complete, combined thoracolumbar approach with decompression of spinal cord, cauda equina or nerve root(s), lower thoracic or lumbar; single segment
63088	Vertebral corpectomy (vertebral body resection), partial or complete, combined thoracolumbar approach with decompression of spinal cord, cauda equina or nerve root(s), lower thoracic or lumbar; each additional segment (List separately in addition to code for primary procedure)
63090	Vertebral corpectomy (vertebral body resection), partial or complete, transperitoneal or retroperitoneal approach with decompression of spinal cord, cauda equina or nerve root(s), lower thoracic, lumbar, or sacral; single segment
63091	Vertebral corpectomy (vertebral body resection), partial or complete, transperitoneal or retroperitoneal approach with decompression of spinal cord, cauda equina or nerve root(s), lower thoracic, lumbar, or sacral; each additional segment (List separately in addition to code for primary procedure)
63101	Vertebral corpectomy (vertebral body resection), partial or complete, lateral extracavitary approach with decompression of spinal cord and/or nerve root(s) (e.g., for tumor or retropulsed bone fragments); thoracic, single segment
63102	Vertebral corpectomy (vertebral body resection), partial or complete, lateral extracavitary approach with decompression of spinal cord and/or nerve root(s) (e.g., for tumor or retropulsed bone fragments); lumbar, single segment

CPT Code	Description
63103	Vertebral corpectomy (vertebral body resection), partial or complete, lateral extracavitary approach with decompression of spinal cord and/or nerve root(s) (e.g., for tumor or retropulsed bone fragments); thoracic or lumbar, each additional segment (List separately in addition to code for primary procedure)
63170	Laminectomy with myelotomy (e.g., Bischof or DREZ type), cervical, thoracic, or thoracolumbar
63172	Laminectomy with drainage of intramedullary cyst/syrinx; to subarachnoid space
63173	Laminectomy with drainage of intramedullary cyst/syrinx; to peritoneal or pleural space
63185	Laminectomy with rhizotomy; 1 or 2 segments
63190	Laminectomy with rhizotomy; more than 2 segments
63191	Laminectomy with section of spinal accessory nerve
63197	Laminectomy with cordotomy, with section of both spinothalamic tracts, 1 stage, thoracic
63200	Laminectomy, with release of tethered spinal cord, lumbar
63250	Laminectomy for excision or occlusion of arteriovenous malformation of spinal cord; cervical
63251	Laminectomy for excision or occlusion of arteriovenous malformation of spinal cord; thoracic
63252	Laminectomy for excision or occlusion of arteriovenous malformation of spinal cord; thoracolumbar
63265	Laminectomy for excision or evacuation of intraspinal lesion other than neoplasm, extradural; cervical
63266	Laminectomy for excision or evacuation of intraspinal lesion other than neoplasm, extradural; thoracic
63267	Laminectomy for excision or evacuation of intraspinal lesion other than neoplasm, extradural; lumbar
63270	Laminectomy for excision of intraspinal lesion other than neoplasm, intradural; cervical
63271	Laminectomy for excision of intraspinal lesion other than neoplasm, intradural; thoracic
63272	Laminectomy for excision of intraspinal lesion other than neoplasm, intradural; lumbar
63275	Laminectomy for biopsy/excision of intraspinal neoplasm; extradural, cervical
63277	Laminectomy for biopsy/excision of intraspinal neoplasm; extradural, lumbar
63280	Laminectomy for biopsy/excision of intraspinal neoplasm; intradural, extramedullary, cervical
63282	Laminectomy for biopsy/excision of intraspinal neoplasm; intradural, extramedullary, lumbar
63285	Laminectomy for biopsy/excision of intraspinal neoplasm; intradural, intramedullary, cervical
63286	Laminectomy for biopsy/excision of intraspinal neoplasm; intradural, intramedullary, thoracic
63287	Laminectomy for biopsy/excision of intraspinal neoplasm; intradural, intramedullary, thoracolumbar
63290	Laminectomy for biopsy/excision of intraspinal neoplasm; combined extradural-intradural lesion, any level
63300	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; extradural, cervical
63301	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; extradural, thoracic by transthoracic approach
63302	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; extradural, thoracic by thoracolumbar approach
63303	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; extradural, lumbar or sacral by transperitoneal or retroperitoneal approach
63304	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; intradural, cervical
63305	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; intradural, thoracic by transthoracic approach
63306	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; intradural, thoracic by thoracolumbar approach
63307	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; intradural, lumbar or sacral by transperitoneal or retroperitoneal approach

CPT Code	Description
63308	Vertebral corpectomy (vertebral body resection), partial or complete, for excision of intraspinal lesion, single segment; each additional segment (List separately in addition to codes for single segment)

CPT® is a registered trademark of the American Medical Association

Description of Services

Lumbar Spinal Stenosis (LSS) is a narrowing of the spinal canal that compresses the neural elements in the lower back. It may be caused by trauma, tumor, infection, or congenital defects but is predominately caused by degenerative changes in the intervertebral discs and the ligaments and bone structures of the spine. These changes typically begin with a breakdown of the discs with consequent collapse of disc space, which leads to disc bulge and herniation, and transference of weight to the facet joints. This in turn leads to cartilage erosion and compensatory growth of new bone (bone spurs) over the facet joints as well as thickening of ligaments around the facet joints to help support the vertebrae. Surgery may be performed if symptoms do not respond adequately to nonsurgical approaches and continue to cause poor quality of life (AAOS, 2013).

First-line treatments for symptomatic Lumbar Spinal Stenosis include rest, NSAIDs, muscle relaxants, corset use, physical therapy, and lumbar epidural steroid injections. For persons with moderate to severe symptoms, surgical decompression with or without Spinal Fusion and discectomy may be indicated but are associated with serious complications and high operative risk, particularly for elderly patients. The effectiveness of nonsurgical treatments, the extent of pain, and patient preferences may all factor into the decision to have surgery [National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), 2016].

Arthrodesis (fusion) procedures in the lumbar (lower) spine are surgical procedures that join two or more lumbar vertebrae together into one solid bony structure. These procedures may be used to treat spine instability, cord compression due to severe degenerative disc disease, fractures in the lumbar spine, or destruction of the vertebrae by infection or tumor. There are several methods or approaches to this surgery.

The most common approach to Arthrodesis (fusion) of the lumbar spine is the posterior approach. After the vertebrae are exposed through the back, pressure on the nerve roots and/or spinal cord is removed (“decompressed”). This usually includes removing part or all of the nearby lamina bone, facet joints, any free disc fragments, or filing down any nearby bone spurs to relieve the nerves inside the spinal canal of tension and pressure.

In preparation for the Spinal Fusion, a layer of bone off the back surfaces of the affected spinal column is removed. Small strips of bone called bone grafts are then removed from the top rim of the pelvis and placed over the now exposed bone surfaces of the spinal column. As healing occurs, the bone strips will fuse across the spaces in between the vertebral bodies, such as the disc spaces or the facet joint spaces. To reinforce the fusion procedure, the bones may be fixated in place using a combination of metal screws, rods, and plates. This instrumentation holds together the vertebrae to be fused, to prevent them from moving during the bone healing process.

Clinical Evidence

Dynamic Stabilization Systems

Due to the lack of data from well-designed, long-term, randomized controlled clinical trials, current evidence is insufficient to permit conclusions about whether any beneficial effect from dynamic stabilization provides a significant advantage over conventional fusion techniques. The published evidence is not robust; a majority of the studies are retrospective or prospective case series and lack controls. In addition, the complication rates and reoperation rates for dynamic stabilization compared with conventional fusion are unknown.

In 2024, Nassr et al. performed a multicenter randomized controlled trial (RCT) of 321 patients with lumbar spinal stenosis and grade-I degenerative spondylolisthesis. A total of 321 adult patients were randomized in a 2:1 fashion, with 219 patients assigned to undergo facet arthroplasty and 102 patients assigned to undergo fusion. Of these, 113 patients (51.6%) in the arthroplasty group and 47 (46.1%) in the fusion group who had either reached 24 months of postoperative follow-up or were deemed early clinical failures were included in the primary outcome analysis. The arthroplasty group had a higher proportion of patients who achieved composite clinical success than did the fusion group (73.5% versus 25.5%; $p < 0.001$), equating to a between-group difference of 47.9% (95% confidence interval, 33.0% to 62.8%). The arthroplasty group outperformed the fusion group in most patient-reported outcome measures (including the ODI, VAS back pain, and all ZCQ component scores) at 24 months postoperatively. There were no significant differences between

groups in surgical variables or complications, except that the fusion group had a higher rate of developing symptomatic adjacent segment degeneration. The study demonstrated that decompression plus lumbar facet arthroplasty was associated with superior PROMs across multiple metrics, lower rates of new or progressive neurologic symptoms, and lower rates of symptomatic adjacent segment degeneration, equating to higher rates of composite clinical success, compared with decompression plus fusion, at 24 months postoperatively. Long-term follow-up will be necessary to determine differences in implant longevity, PROMs, and radiographic parameters such as stability of the spondylolisthesis and maintenance of motion beyond 2 years. A future randomized controlled trial may be considered to compare lumbar facet arthroplasty versus decompression alone in a broader sample of patients. The primary limitation of this study is the relatively short postoperative follow-up, which precludes evaluation of the long-term durability of lumbar facet arthroplasty. A second limitation is that industry funding was utilized to perform this study. Third, this study was unable to blind surgeons, patients, or radiologists to the patients' treatment allocation postoperatively. Therefore, detection bias is a distinct possibility. Fourth, the trial utilized strict inclusion and exclusion criteria to mitigate the impact of confounding variables on the outcomes reported. Finally, this study reports the primary outcome in only approximately one half of the randomized sample; however, this is consistent with both the predetermined statistical plan and previous randomized controlled trials when there is a sufficiently large between-group difference at a preplanned interim analysis.

In 2022, Pinter et al. conducted an interim analysis on the one-year safety profile and clinical and radiographic outcomes of 153 participants randomized to the investigational arm of the FDA investigational device exemption (IDE) clinical trial for the Total Posterior Spine System (TOPS™) (Premia Spine USA, Norwalk, CT) device. Among the participants, 145 devices were implanted at L4-5 and 8 at L3-4. 105 participants had reached the one year follow up and are included in the results. The safety profile showed 11 total complications and included new neurological deficits, dural tears infection, seroma, hematoma as well as retained drains, misplaced pedicle screws and screw loosening. Nine of these required a total of 13 reoperations. Patient reported outcome measures (PROMs) showed sustained improvement from 6 weeks to 12 months in ODI scores, as well as the mean VAS scores for low back and leg pain. Zurich Claudication Questionnaire (ZCQ) symptom scores also improved. Radiographic parameters included global lordosis, disc height at the index level, and disc angle and spondylolisthesis at the index level and the levels above and below the index level and was evaluated in 90 of the participants. Static radiographic parameters demonstrated increased index disc angle and disc height with a reduction in the magnitude of spondylolisthesis. Comparison of dynamic radiographic parameters showed increased flexion/extension ROM and translation. This analysis reports the early outcomes using a device with FDA IDE. Further information on this ongoing clinical trial can be found at:

<https://www.clinicaltrials.gov/ct2/show/NCT03012776?term=Total+Posterior+Spine+System&draw=2&rank=1>.

Pham et al. (2016) conducted a review of the literature to explore complications associated with the Dynesys stabilization system. The researchers evaluated 21 studies which included a total of 1,166 subjects with a mean age of 55.5 years and a mean follow-up period of 33.7 months. The data demonstrated a surgical-site infection rate of 4.3%, a pedicle screw loosening rate of 11.7%, a pedicle screw fracture rate of 1.6%, and an adjacent-segment disease (ASD) rate of 7.0%. Of studies reporting surgical revision rates, 11.3% of subjects required reoperation. Of subjects who developed ASD, 40.6% required a reoperation for treatment. The authors concluded that the Dynesys stabilization system has a similar complication rate compared with lumbar fusion studies and has a slightly lower incidence of ASD.

Clinical Practice Guidelines

North American Spine Society (NASS)

A 2014 (revised in 2019) NASS coverage policy recommendation, entitled *Interspinous Fixation with Fusion* which addresses “flexible fusion,” which is defined as dynamic stabilization without arthrodesis for the treatment of degenerative lumbar spondylolisthesis. Due to the paucity of literature addressing the outcomes of these procedures, the workgroup was unable to make a recommendation. For future research, the workgroup recommended development of a large multicenter registry database, as well as prospective studies, with long-term follow-up comparing flexible fusion to medical or interventional treatment of this condition.

Isolated Facet Fusion

Evidence is limited to small, uncontrolled trials with lack of blinding or long-term follow-up. Randomized, controlled trials comparing these allograft materials to standardized autograft materials are needed to determine long-term efficacy and impact on health outcomes. No studies were found that discussed facet fusion when done alone without an accompanying decompressive procedure. The current published evidence is insufficient to determine whether isolated facet arthroplasty is as effective or as safe as spinal fusion, which is the current standard for surgical treatment of degenerative disc disease.

Gavaskar and Achimuthu (2010) conducted a prospective study of 30 patients with low-grade degenerative spondylolisthesis of the lumbar and lumbosacral spine who underwent facet fusion using 2 cortical screws and local

cancellous bone grafts. Visual analog scale and Oswestry disability assessment were used to measure outcomes which showed significant improvement at 1-year follow-up. The authors found that patients with degenerative spondylolisthesis with lower grade slips and normal anterior structures represent an ideal indication for facet fusion. The study is limited by short term follow-up, subjective outcomes and lack of comparison to other treatment modalities.

Multiple Serial/Staged Spine Procedures

There is insufficient evidence of efficacy to support dividing spine decompression procedures into serial, multiple, or staged sessions when one session can address all sites.

Total Facet Arthroplasty

Due to the lack of data from well-designed, long-term, randomized controlled clinical trials, current evidence is insufficient to permit conclusions about benefits and safety of facet arthroplasty. For individuals who have lumbar spinal stenosis who receive spinal decompression with facet arthroplasty, the evidence includes a preliminary report of a randomized controlled trial. Interim results from a pivotal trial of the ACADIA Facet Replacement System were reported in 2012. No additional publications from this trial, which was expected to be completed October 2015, have been identified to date.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Spinal Fusion Devices

Products used for spinal fusion and decompression devices are extensive. Refer to the following website for more information and search by product name in device name section:

<http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnm.cfm>. (Accessed October 23, 2024)

Dynamic Stabilization Systems

On June 15, 2023, the FDA granted premarket approval (PMA) of the TOPS™ System (Premia Spine USA, Norwalk, CT), a motion preserving spinal implant intended to stabilize the spine following decompression without using rigid fixation.

Further information can be found at: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P220002>. (Accessed October 23, 2024)

Facet Arthroplasty

No facet arthroplasty devices have been approved by the U.S. Food and Drug Administration (FDA) at this time.

References

American Academy of Orthopaedic Surgeons (AAOS). Lumbar Spinal Stenosis. December 2013.

Chen YN, Yang SH, Chou SC, et al. The role of sacral laminoplasty in the management of spina bifida and sacral cystic lesions: case series. *Neurosurg Focus*. 2019 Oct 1;47(4):E20.

Gavaskar AS, Achimuthu R. Transfacetal fusion for low-grade degenerative spondylolisthesis of the lumbar spine: results of a prospective single center study. *J Spinal Disord Tech*. 2010 May;23(3):162-5.

Gellhorn AC, Katz JN, Suri P. Osteoarthritis of the spine: the facet joints. *Nat Rev Rheumatol*. 2013 Apr;9(4):216-24.

McAfee P, Khoo LT, Pimenta L, et al. Treatment of lumbar spinal stenosis with a total posterior arthroplasty prosthesis: implant description, surgical technique, and a prospective report on 29 patients. *Neurosurg Focus*. 2007;22(1):E13.

Mendenhall S, Mobasser D, Relyea K, et al. Spinal instrumentation in infants, children, and adolescents: a review. *J Neurosurg Pediatr*. 2019 Jan 1;23(1):1-15.

Nassar A, Coric D, Pinter Z, et al. Lumbar Facet Arthroplasty Versus Fusion for Grade-I Degenerative Spondylolisthesis with Stenosis: A Prospective Randomized Controlled Trial. *The Journal of Bone and Joint Surgery* 106(12):p 1041-1053, June 19, 2024.

Negrini et al. 2016 SOSORT guidelines: orthopaedic and rehabilitation treatment of idiopathic scoliosis during growth. *Scoliosis Spinal Disord* 2018.

North American Spine Society (NASS). Coverage Policy Recommendations: Interspinous Fixation with Fusion. 2014. Revised 2019.

North American Spine Society (NASS). Coverage Policy Recommendations. Lumbar Fusion. 2014a.

Pham M, Mehta V, Patel N, et.al. Complications associated with the Dynesys dynamic stabilization system: a comprehensive review of the literature. Neurosurg Focus. 2016.

Pinter ZW, Freedman BA, Nassr A, et al. A prospective study of lumbar facet arthroplasty in the treatment of degenerative spondylolisthesis and stenosis: Results from the Total Posterior Spine System (TOPS) IDE Study. Clin Spine Surg. 2022 Aug 3.

Qassem, Amir, et al. Noninvasive treatments for acute, subacute and chronic low back pain: A clinical practice guideline from the American College of Physicians. Annals of Internal Medicine. April 2017.

Summers, Jeffrey. International Spine Intervention Society Recommendations for treatment of Cervical and Lumbar Spine Pain. 2013.

Sun S, Li Y, Wang X, et al. Safety and efficacy of laminoplasty versus laminectomy in the treatment of spinal cord tumors: A systematic review and meta-analysis. World Neurosurg. 2019 May;125:136-145.

Veritas Health Inc. Spine-health. Available at: <https://www.spine-health.com/author/veritas-health>. Accessed October 23, 2024.

Policy History/Revision Information

Date	Summary of Changes
06/01/2025	<ul style="list-style-type: none">New Medical Policy

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state, or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state, or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state, or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state, or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.